

# MA3204 - Exercise 7

Throughout the following exercises  $\mathcal{A}$  denotes an abelian category, and  $\mathcal{C}$  denotes an arbitrary category. For the exercises about localizations we will ignore set-theoretical issues.

- Let  $\mathcal{A}$  be an additive category, let  $X \in \mathcal{A}$ , and let  $A^\bullet = (\dots \xrightarrow{d^{-2}} A^{-1} \xrightarrow{d^{-1}} A^0 \xrightarrow{d^0} A^1 \xrightarrow{d^1} \dots)$  be a complex in  $\mathcal{A}$ . Define the complex

$$\mathrm{Hom}_{\mathcal{A}}(A^\bullet, X) = (\dots \xrightarrow{-\circ d^1} \mathrm{Hom}_{\mathcal{A}}(A^1, X) \xrightarrow{-\circ d^0} \mathrm{Hom}_{\mathcal{A}}(A^0, X) \xrightarrow{-\circ d^{-1}} \mathrm{Hom}_{\mathcal{A}}(A^{-1}, X) \xrightarrow{-\circ d^{-2}} \dots)$$

Show that

$$\begin{aligned} Z^n \mathrm{Hom}_{\mathcal{A}}(A^\bullet, X) &= \mathrm{Hom}_{\mathrm{Ch}(\mathcal{A})}(A^\bullet, X[n]) \\ H^n \mathrm{Hom}_{\mathcal{A}}(A^\bullet, X) &= \mathrm{Hom}_{\mathbf{K}(\mathcal{A})}(A^\bullet, X[n]) \end{aligned}$$

- Let  $\mathcal{C}$  be a category, and let  $S \subset \mathrm{Mor} \mathcal{C}$  be a class of morphisms in  $\mathcal{C}$ . Consider the localization  $\mathcal{C}[S^{-1}]$  defined in the lecture, whose morphism spaces consists of strings of morphisms in  $\mathcal{C}$  and formal inverses of morphisms in  $S$ .
  - Show that  $\mathcal{C}[S^{-1}]$  is a category, with identity morphisms given by the empty strings  $[X, \emptyset, X]$ .
  - Show that we have a functor  $Q: \mathcal{C} \rightarrow \mathcal{C}[S^{-1}]$  given by the identity on objects, and sending a morphism  $f: X \rightarrow Y$  to the string  $Q(f) = [X, f, Y]$ . Furthermore, show that if  $s \in S$ , then  $Q(s)$  is an isomorphism
  - Show that  $(\mathcal{C}[S^{-1}], Q)$  satisfies the universal property of the localization. In other words, if  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a functor satisfying that  $F(s)$  is an isomorphism for all  $s \in S$ , then show that there exists a unique functor

$$\bar{F}: \mathcal{C}[S^{-1}] \rightarrow \mathcal{D}$$

satisfying  $F = \bar{F} \circ Q$ .


- Let  $\mathcal{C}$  be a category, and let  $S \subset \mathrm{Mor} \mathcal{C}$  be a class of morphisms in  $\mathcal{C}$ . Assume  $S$  admits a calculus of right fractions. We defined two right fractions  $(s_1, \alpha_1)$  and  $(s_2, \alpha_2)$  from  $X$  to  $Y$  to be equivalent, written  $(s_1, \alpha_1) \sim (s_2, \alpha_2)$ , if there exists a right fraction  $(s_3, \alpha_3)$  and a commutative diagram

$$\begin{array}{ccccc} & & X_1 & & \\ & \swarrow & \uparrow & \searrow & \\ X & \xleftarrow{s_1} & X_3 & \xrightarrow{\alpha_3} & Y \\ & \swarrow & \downarrow & \searrow & \\ & & X_2 & & \end{array}$$

Show that  $\sim$  is an equivalence relation on the class of right fractions from  $X$  to  $Y$ . We denote the equivalence class of  $(s, \alpha)$  by  $[s, \alpha]$ .

- Let  $\mathcal{C}$  be a category, and let  $S \subset \mathrm{Mor} \mathcal{C}$  be a class of morphisms in  $\mathcal{C}$ . Assume  $S$  admits a calculus of right fractions. Recall that a composite of two right fractions  $X \xleftarrow{s_1} X_1 \xrightarrow{\alpha_1} Y$  and  $Y \xleftarrow{t_1} Y_1 \xrightarrow{\beta_1} Z$  is a right fraction  $(s_1 \circ t'_1, \beta_1 \circ \alpha'_1)$  where  $t'_1 \in S$  and where we have commutative

square



obtained from axiom (RF2).

(a) Show that the equivalence class of  $(s_1 \circ t'_1, \beta_1 \circ \alpha'_1)$  is independent of the choice of morphisms  $t'_1$  and  $\alpha'_1$  above. Let  $(t_1, \beta_1) \circ (s_1, \alpha_1)$  denote this unique class.

(b) Show that if  $(t_1, \beta_1) \sim (t_2, \beta_2)$ , then

$$(t_1, \beta_1) \circ (s_1, \alpha_1) = (t_2, \beta_2) \circ (s_1, \alpha_1)$$

(c) Show that if  $(s_1, \alpha_1) \sim (s_2, \alpha_2)$ , then

$$(t_1, \beta_1) \circ (s_1, \alpha_1) = (t_1, \beta_1) \circ (s_2, \alpha_2)$$

(d) Consider  $S^{-1}\mathcal{C}$  as defined in the lecture, whose morphisms spaces are equivalence classes of right fractions. Show that the composition law above induces a well-defined map

$$\circ: \text{Hom}_{S^{-1}\mathcal{C}}(Y, Z) \times \text{Hom}_{S^{-1}\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{S^{-1}\mathcal{C}}(X, Z).$$

5. Let  $\mathcal{C}$  be a category, and let  $S \subset \text{Mor } \mathcal{C}$  be a class of morphisms in  $\mathcal{C}$ . Assume  $S$  admits a calculus of right fractions. Using the previous exercises, show that  $S^{-1}\mathcal{C}$  is a category, with identity morphisms given by the equivalence classes of the right fractions  $X \xleftarrow{1} X \xrightarrow{1} X$ .

6. Let  $\mathcal{C}$  be a category, and let  $S \subset \text{Mor } \mathcal{C}$  be a class of morphisms in  $\mathcal{C}$ . Assume  $S$  admits a calculus of right fractions. Show that we have an *isomorphism* of categories

$$F: S^{-1}\mathcal{C} \rightarrow \mathcal{C}[S^{-1}]$$

given by the identity on objects, and sending a right fraction  $X \xleftarrow{s} Y' \xrightarrow{\alpha} Y$  to the string  $[X, \alpha, s^-, Y]$ .

7. Let  $\mathcal{B}$  be an additive category, and let  $S \subset \text{Mor } \mathcal{B}$  be a class of morphisms in  $\mathcal{C}$ . Assume  $S$  admits a calculus of right fractions. Let  $[s_1, \alpha_1]$  and  $[s_2, \alpha_2]$  be two morphisms in  $\text{Hom}_{S^{-1}\mathcal{B}}(X, Y)$ . In the lecture we explained how one can find a morphism  $s \in S$  and morphisms  $\beta_1, \beta_2$  in  $\mathcal{B}$  such that  $[s_1, \alpha_1] = [s, \beta_1]$  and  $[s_2, \alpha_2] = [s, \beta_2]$ . We then defined

$$[s_1, \alpha_1] + [s_2, \alpha_2] = [s, \beta_1 + \beta_2].$$

The goal of this exercise is to show that this operation is well-defined, i.e. that if  $[s_1, \alpha_1] = [t, \beta'_1]$  and  $[s_2, \alpha_2] = [t, \beta'_2]$ , then

$$[s, \beta_1 + \beta_2] = [t, \beta'_1 + \beta'_2].$$

In the following  $[s, \alpha], [s, \alpha'], [s, \beta], [s, \beta']$  are morphisms in  $\text{Hom}_{S^{-1}\mathcal{B}}(X, Y)$

(a) Show that  $[s, \alpha] = [s, \alpha']$  if and only if there exists some  $t \in S$  such that  $\alpha \circ t = \alpha' \circ t$  and  $s \circ t \in S$ .

(b) Show that if  $[s, \alpha] = [s, \alpha']$  and  $[s, \beta] = [s, \beta']$ , then

$$[s, \alpha + \beta] = [s, \alpha' + \beta'].$$

(c) Now assume  $s, s_1, s_2, t$  and  $\alpha_1, \alpha_2, \beta_1, \beta_2, \beta'_1, \beta'_2$  are as above. Applying (RF2) to  $X_1 \xrightarrow{s} X \xleftarrow{t} X'_1$ , show that we can find  $\tilde{s} \in S$  and morphisms  $\gamma_1, \gamma_2, \gamma'_1, \gamma'_2$  such that

$$\begin{array}{lll}
 [s, \beta_1] = [\tilde{s}, \gamma_1] & [s, \beta_2] = [\tilde{s}, \gamma_2] & [s, \beta_1 + \beta_2] = [\tilde{s}, \gamma_1 + \gamma_2] \\
 [t, \beta'_1] = [\tilde{s}, \gamma'_1] & [t, \beta'_2] = [\tilde{s}, \gamma'_2] & [t, \beta'_1 + \beta'_2] = [\tilde{s}, \gamma'_1 + \gamma'_2].
 \end{array}$$

- (d) Using part (b) and (c), conclude that  $[s, \beta_1 + \beta_2] = [t, \beta'_1 + \beta'_2]$ .
8. Let  $\mathcal{B}$  be an additive category, and let  $S \subset \text{Mor } \mathcal{B}$  be a class of morphisms in  $\mathcal{C}$ . Assume  $S$  admits a calculus of right fractions. The goal of this exercise is to fill in the remaining details of the proof that  $S^{-1}\mathcal{B}$  is preadditive.
- (a) Show that the operation  $+$  defined in the previous exercise makes  $\text{Hom}_{S^{-1}\mathcal{B}}(X, Y)$  into an abelian group with zero element  $[1_X, 0]$ .
- (b) Show that composition in  $S^{-1}\mathcal{B}$  is bilinear with respect to  $+$ .
9. Let  $X^\bullet \in \mathbf{K}(\mathcal{A})$  be a complex. Show that  $X^\bullet \cong 0$  in  $\mathbf{D}(\mathcal{A})$  if and only if  $X^\bullet$  is exact.
10. Let  $f : A^\bullet \rightarrow B^\bullet$  be a morphism in  $\mathbf{K}(\mathcal{A})$ . Show that the following are equivalent.
- (a)  $f$  factors through an exact complex.
- (b) There exists a quasi-isomorphism  $q : H^\bullet \rightarrow A^\bullet$  such that  $f \circ q = 0$ .
- (c)  $f \cdot \text{id}_{A^\bullet}^{-1} = 0$  in  $\mathbf{D}(\mathcal{A})$ .