



Please justify your answers! Note that *how* you arrive at an answer is more important than the answer itself.

- 1] Let T be a linear operator on the space of polynomials \mathcal{P}_2 of degree at most 2 defined by $Tf(x) = -f(x) - f'(x)$.
- Find the matrix representation of T with respect to the basis $1, x, x^2$ of \mathcal{P}_2 and its characteristic polynomial.
 - Find the eigenvalue(s) and eigenvector(s) of T .

- 2] Let $A \in \mathcal{M}_{m \times n}(\mathbb{C})$ and $B \in \mathcal{M}_{n \times m}(\mathbb{C})$, and let $\lambda \in \mathbb{C}$ be any nonzero scalar. Show that λ is an eigenvalue of AB if and only if λ is an eigenvalue of BA .

- 3] Let $A \in \mathcal{M}_{n \times n}(\mathbb{C})$ be a normal matrix. Prove that

$$\det(A) = \prod_{j=1}^n \lambda_j,$$

where the λ_j 's are the (not necessarily distinct) eigenvalues of A .

- 4] (*Exam 2017, Problem 1a*)

- a) Find the singular value decomposition for the matrix

$$A = \begin{bmatrix} 1 & 1 & -1 \\ 1 & 1 & -1 \end{bmatrix}.$$

- b) The linear system

$$\begin{aligned} x_1 + x_2 - x_3 &= 1 \\ x_1 + x_2 - x_3 &= 1 \end{aligned}$$

has infinitely many solutions. Find the solution with minimal Euclidean norm $\|\cdot\|_2$.

c) The linear system

$$x_1 + x_2 - x_3 = 1$$

$$x_1 + x_2 - x_3 = 2$$

is inconsistent, and has no solution. Find the unique best approximation to a solution having minimum norm.

d) Prove that an $(n \times n)$ matrix A of full rank has a polar decomposition using the singular value decomposition of A . Hence, show that there exists an $(n \times n)$ unitary matrix W and a positive definite (not just semi-definite) $(n \times n)$ matrix P such that $A = WP$.