

## Inner products

Definition: An *inner product* on a real vector space  $V$  is an operation (function) that assigns to each pair of vectors  $(\vec{u}, \vec{v})$  in  $V$  a **scalar**  $\langle \vec{u}, \vec{v} \rangle$  satisfying the following axioms:

1.  $\langle \vec{u}, \vec{v} \rangle = \langle \vec{v}, \vec{u} \rangle$  (Symmetry)
2.  $\langle \vec{u} + \vec{v}, \vec{w} \rangle = \langle \vec{u}, \vec{w} \rangle + \langle \vec{v}, \vec{w} \rangle$  (Additivity)
3.  $\langle k \vec{u}, \vec{v} \rangle = k \langle \vec{u}, \vec{v} \rangle$  (Homogeneity)
4.  $\langle \vec{v}, \vec{v} \rangle \geq 0$  and  $\langle \vec{v}, \vec{v} \rangle = 0$  iff  $\vec{v} = \vec{0}$  (Positivity)

Theorem (basic properties): Given vectors  $\vec{u}, \vec{v}, \vec{w}$  in an inner product space  $V$ , and a scalar  $k$ , the following properties hold:

- $\langle \vec{0}, \vec{v} \rangle = \langle \vec{v}, \vec{0} \rangle = 0$
- $\langle \vec{u} - \vec{v}, \vec{w} \rangle = \langle \vec{u}, \vec{w} \rangle - \langle \vec{v}, \vec{w} \rangle$
- $\langle \vec{u}, \vec{v} + \vec{w} \rangle = \langle \vec{u}, \vec{v} \rangle + \langle \vec{u}, \vec{w} \rangle$
- $\langle \vec{u}, \vec{v} - \vec{w} \rangle = \langle \vec{u}, \vec{v} \rangle - \langle \vec{u}, \vec{w} \rangle$
- $\langle \vec{u}, k\vec{v} \rangle = k \langle \vec{u}, \vec{v} \rangle$

## Norm and distance in an inner product space

Definition: If  $V$  is a real inner product space then we define

- The norm (or length) of  $\vec{v}$ :

$$\|\vec{v}\| = \sqrt{\langle \vec{v}, \vec{v} \rangle}$$

- The distance between  $\vec{u}$  and  $\vec{v}$ :

$$d(\vec{u}, \vec{v}) = \|\vec{u} - \vec{v}\| = \sqrt{\langle \vec{u} - \vec{v}, \vec{u} - \vec{v} \rangle}$$

Theorem (basic properties): Given vectors  $\vec{u}, \vec{v}$  in an inner product space  $V$ , and a scalar  $k$ , the following properties hold:

- $\|\vec{v}\| \geq 0$  and  $\|\vec{v}\| = 0$  iff  $\vec{v} = \vec{0}$ .
- $\|k\vec{v}\| = |k| \|\vec{v}\|$
- $d(\vec{u}, \vec{v}) = d(\vec{v}, \vec{u})$
- $d(\vec{u}, \vec{v}) \geq 0$  and  $d(\vec{u}, \vec{v}) = 0$  iff  $\vec{u} = \vec{v}$ .

## Angle between vectors

Theorem (Cauchy-Schwarz): If  $u$  and  $v$  are vectors in an inner vector space, then

$$|\langle u, v \rangle| \leq \|u\| \|v\|$$

Definition: The angle between two vectors  $u$  and  $v$  in an inner vector space is defined as

$$\theta = \cos^{-1} \frac{\langle u, v \rangle}{\|u\| \|v\|}$$

Theorem (the triangle inequality): If  $u, v$  and  $w$  are vectors in an inner vector space, then

- $\|u + v\| \leq \|u\| + \|v\|$
- $d(u, v) \leq d(u, w) + d(w, v)$

## Orthogonality

Definition: Two vectors  $u$  and  $v$  in an inner vector space are called *orthogonal* if  $\langle u, v \rangle = 0$ .

Clearly  $u \perp v$  iff the angle between them is  $\theta = \frac{\pi}{2}$ .

Theorem (the Pythagorean theorem): If  $u$  and  $v$  are **orthogonal** vectors in an inner vector space, then

$$\|u + v\|^2 = \|u\|^2 + \|v\|^2$$

Definition: Let  $W$  be a subspace of an inner product space  $V$ . The set of vectors in  $V$  which are orthogonal to **every** vector in  $W$  is called the *orthogonal complement* of  $W$  and it is denoted by  $W^\perp$ .

Theorem: The orthogonal complement has the following properties:

- $W^\perp$  is a subspace of  $V$ .
- $W \cap W^\perp = \{\vec{0}\}$ .
- If  $V$  has finite dimension then  $(W^\perp)^\perp = W$ .

## Orthogonal sets, orthonormal sets

Let  $(V, \langle \cdot | \cdot \rangle)$  be an inner product space and let  $S$  be a set of vectors in  $V$ .

Definition: The set  $S$  is called *orthogonal* if any two vectors in  $S$  are orthogonal.

The set  $S$  is called *orthonormal* if it is orthogonal and any vector in  $S$  has norm 1.

Theorem: Every orthogonal set of nonzero vectors is linearly independent.

Definition: A set of vectors  $S$  is called an *orthogonal basis* (OGB) for  $V$  if  $S$  is a basis and an orthogonal set (that is,  $S$  is a basis where all vectors are perpendicular).

A set of vectors  $S$  is called an *orthonormal basis* (ONB) for  $V$  if  $S$  is a basis and an orthonormal set (that is,  $S$  is a basis where all vectors are perpendicular and have norm 1).

## Orthogonal sets, orthonormal sets

Let  $(V, \langle \cdot | \cdot \rangle)$  be an inner product space.

Theorem: If  $S = \{v_1, v_2, \dots, v_n\}$  is an orthogonal basis in  $V$  and  $u$  is any vector in  $V$ , then

$$u = \frac{\langle u, v_1 \rangle}{\|v_1\|^2} v_1 + \frac{\langle u, v_2 \rangle}{\|v_2\|^2} v_2 + \dots + \frac{\langle u, v_n \rangle}{\|v_n\|^2} v_n$$

If  $S = \{v_1, v_2, \dots, v_n\}$  is an orthonormal basis in  $V$  and  $u$  is any vector in  $V$ , then

$$u = \langle u, v_1 \rangle v_1 + \langle u, v_2 \rangle v_2 + \dots + \langle u, v_n \rangle v_n$$

## Gram-Schmidt process

Theorem: Every nonzero finite dimensional inner product space has an orthonormal basis.

Given a basis  $\{u_1, u_2, \dots, u_n\}$ , to find an orthogonal basis  $\{v_1, v_2, \dots, v_n\}$  we use the following procedure:

Step 1.  $v_1 = u_1$

Step 2.  $v_2 = u_2 - \frac{\langle u_2, v_1 \rangle}{\|v_1\|^2} v_1$

Step 3.  $v_3 = u_3 - \frac{\langle u_3, v_1 \rangle}{\|v_1\|^2} v_1 - \frac{\langle u_3, v_2 \rangle}{\|v_2\|^2} v_2$

Step 4.  $v_4 = u_4 - \frac{\langle u_4, v_1 \rangle}{\|v_1\|^2} v_1 - \frac{\langle u_4, v_2 \rangle}{\|v_2\|^2} v_2 - \frac{\langle u_4, v_3 \rangle}{\|v_3\|^2} v_3$

and so on for  $n$  steps, where  $n = \dim(V)$ .

To obtain an orthonormal basis, we simply normalize the orthogonal basis obtained above.