



Norwegian University of Science
and Technology
Department of Mathematics

MA2501: Numerical Methods
Spring 2016

Solutions to exercise set 7

1 Cf. Cheney and Kincaid, Exercise 4.1.9

Consider the data points

x_i	0	1	2	4	6
$f(x_i)$	1	9	23	93	259

- Find the interpolation polynomial through these data points using Newton interpolation, and compute an approximation of f at $x = 3$.
- Do the same using only the first four interpolation points.

Possible solution:

- a) We obtain the following table for the divided differences:

0	1			
		8		
1	9		3	
		14		1
2	23		7	0
		35		1
4	93		12	
		83		
6	259			

The interpolation polynomial (in nested Newton form) is therefore

$$p(x) = 1 + x(8 + (x - 1)(3 + (x - 2) + 0 \cdot (x - 4))).$$

(Obviously, one can omit the term $0 \cdot (x - 4)$ in that expression). Evaluating this polynomial, we see that $p(3) = 49$.

- b) For the interpolation polynomial of degree three through the first four interpolation points, we can simply reuse the previously computed table of divided differences and read off the result from there using only the first four columns. In this particular case, we obtain the same interpolation polynomial p as before (therefore, obviously, also its value at the point $x = 3$ is the same as above).

2 Suppose we have the nodes

$$x_0 = -2, x_1 = -1, x_2 = 1 \text{ and } x_3 = 4,$$

and know the divided differences

$$f[x_3] = 11, f[x_2, x_3] = 5, f[x_2, x_0, x_1] = -2, \text{ and } f[x_0, x_2, x_1, x_3] = 0.6.$$

What is $f(-1)$?

Possible solution:

To start we exploit the fact that divided differences are invariant under permutations of the nodes. Thus, sorting the nodes in increasing order, we have:

$$f[x_3] = 11, f[x_2, x_3] = 5, f[x_0, x_1, x_2] = -2, \text{ and } f[x_0, x_1, x_2, x_3] = 0.6.$$

We seek $f(-1) = f(x_1) = f[x_1]$. From the recursive definition of divided differences:

$$f[x_0, x_1, x_2, x_3] = \frac{f[x_1, x_2, x_3] - f[x_0, x_1, x_2]}{x_3 - x_0}$$

which rearranged to isolate the unknown $f[x_1, x_2, x_3]$ becomes

$$f[x_1, x_2, x_3] = f[x_0, x_1, x_2] + (x_3 - x_0)f[x_0, x_1, x_2, x_3] = -2 + (4 - (-2))0.6 = 1.6.$$

Similarly we find $f[x_1, x_2]$ as

$$f[x_1, x_2] = f[x_2, x_3] - (x_3 - x_1)f[x_1, x_2, x_3] = 5 - (4 - (-1))1.6 = -3,$$

$f[x_2]$ as

$$f[x_2] = f[x_3] - (x_3 - x_2)f[x_2, x_3] = 11 - (4 - 1)5 = -4,$$

and finally $f[x_1] = f(-1)$ as

$$f[x_1] = f[x_2] - (x_2 - x_1)f[x_1, x_2] = -4 - (1 - (-1))(-3) = 2.$$

- 3 Use the interpolation error formula to find a bound for the error, and compare the bound to the actual error for the case $n = 2$ for Exercise 6 in Exercise set 6.

Possible solution:

- a) For $f(x) = x \ln x$ we first calculate $f'(x) = \ln x + 1$, $f''(x) = 1/x$ and $f^{(3)}(x) = -1/x^2$. Since $|f^{(3)}(x)| = 1/x^2$ is a strictly decreasing function of x for positive x we find a bound M on the interval $(8.3, 8.7)$ for $n = 2$ by $M = 1/8.3^2$. Employing the First Interpolation Error Theorem in the textbook with $n = 2$ and $x = 8.4$ on this interval gives the error bound:

$$\begin{aligned} |f(8.4) - p_2(8.4)| &\leq \frac{1}{(2+1)!} \frac{1}{8.3^2} |(8.4 - 8.3)| |(8.4 - 8.6)| |(8.4 - 8.7)| \\ &= \frac{0.1 \cdot 0.2 \cdot 0.3}{6 \cdot 8.3^2} \leq 1.46 \times 10^{-5}. \end{aligned}$$

The true error is 1.38×10^{-5} correct to the digits given. As expected this is lower than the error bound.

- b) For $f(x) = x^4 - x^3 + x^2 - x + 1$ we again calculate $f'(x) = 4x^3 - 3x^2 + 2x - 1$, $f''(x) = 12x^2 - 6x + 2$ and $f^{(3)}(x) = 24x - 6$. Since $|f^{(3)}(x)| = |24x - 6|$ is easily observed to be a strictly decreasing function of x for negative x , we find a bound M on the interval $(-0.5, 0)$ for $n = 2$ by $M = |f^{(3)}(-0.5)| = 18$. Employing the First Interpolation Error Theorem in the textbook with $n = 2$ and $x = -1/3$ on this interval gives the error bound:

$$\begin{aligned} |f(-1/3) - p_2(-1/3)| &\leq \frac{1}{(2+1)!} 18 |(-1/3 + 1/2)| |(-1/3 + 1/4)| |(-1/3 - 0)| \\ &= \frac{18}{6 \cdot 6 \cdot 12 \cdot 3} = \frac{1}{72} \leq 1.4 \times 10^{-2}. \end{aligned}$$

The true error is 9.6×10^{-3} correct to the digits given. As expected this is lower than the error bound.

- 4 Given the function $f(x) = e^x \sin x$ on the interval $[-4, 2]$.

- a) Show by induction

$$f^{(m)}(x) = \frac{d^m}{dx^m} f(x) = 2^{m/2} e^x \sin(x + m\pi/4).$$

- b) Let $p_n(x)$ be the polynomial that interpolates $f(x)$ in $n+1$ uniformly distributed nodes (including the endpoints). Find an upper bound for the interpolation error on this interval, expressed using n . What must n be to guarantee an error less than 10^{-5} ? Use trial and error or calculate it using MATLAB.
- c) Use MATLAB to verify that the value of n found in **b**), indeed results in an error less than 10^{-5} .

Possible solution:

- a) It is trivial to see that the formula is valid for $m = 0$. Assume now it is correct for some $m \geq 0$. Then

$$\begin{aligned}
 f^{(m+1)}(x) &= \frac{d}{dx} f^{(m)}(x) = 2^{m/2} [e^x \sin(x + m\pi/4) + e^x \cos(x + m\pi/4)] \\
 &= 2^{m/2} e^x [\sin(x + m\pi/4) + \cos(x + m\pi/4)] \\
 &= 2^{m/2} e^x \sqrt{2} \left[\frac{1}{\sqrt{2}} \sin(x + m\pi/4) + \frac{1}{\sqrt{2}} \cos(x + m\pi/4) \right] \\
 &= 2^{m/2} e^x 2^{1/2} [\cos(\pi/4) \sin(x + m\pi/4) + \sin(\pi/4) \cos(x + m\pi/4)] \\
 &= 2^{(m+1)/2} e^x \sin((x + m\pi/4) + \pi/4) \\
 &= 2^{(m+1)/2} e^x \sin(x + (m+1)\pi/4).
 \end{aligned}$$

This shows that the formula holds for $m + 1$ as well, assuming it holds for m . The proof by induction is then complete, and the stated formula is seen to hold for all $m \geq 0$

- b) Our function f has continuous derivatives of any order on the entire real line and on the interval $[-4, 2]$ it satisfies

$$|f^{(n+1)}(x)| = 2^{(n+1)/2} e^x |\sin(x + (n+1)\pi/4)| \leq 2^{(n+1)/2} e^2 = M,$$

where we have used that e^x is a strictly increasing positive function and that $|\sin(x)| \leq 1$ for all x . The Second Interpolation Error Theorem is applicable and we have for $x \in [-4, 2]$

$$\begin{aligned}
 |f(x) - p(x)| &\leq \frac{1}{4(n+1)} M h^{n+1} = \frac{1}{4(n+1)} 2^{(n+1)/2} e^2 \left(\frac{6}{n}\right)^{n+1} \\
 &= \frac{e^2}{4(n+1)} \left(\frac{6\sqrt{2}}{n}\right)^{n+1}.
 \end{aligned}$$

By differentiating, this expression is readily seen to decrease with increasing positive n . Using trial and error we find that the expression has the value 1.27×10^{-5} for $n = 15$ and 2.26×10^{-6} for $n = 16$. Thus $n = 16$ is required to guarantee an accuracy of 10^{-5} .

- c) Using MATLAB we have plotted below $|f(x) - p(x)|$ over the interval, where $p(x)$ is the interpolating polynomial from b) with $n = 16$.

We observe that the error stays several orders of magnitude lower than our requirement of 10^{-5} . This is to be expected. Recall that we were in fact guaranteed an error less than 2.26×10^{-6} , and the theorem only gives fairly loose upper bounds.

Also observe that the errors near the endpoints dominate the errors towards the middle of the interval. This is typical, and illustrates why Chebyshev nodes, which become more clustered as we approach the endpoints, usually gives significantly smaller errors for the same number of nodes.

