

# MA2501, Spring 2019, Numerical Methods

February 11th 2019

## Problem 1

Apply Lagrange interpolation to solve the following problem: find the unique polynomial of degree at most two satisfying

$$p_2(0) = 0, \quad p_2\left(\frac{2}{3}\right) = 1, \quad p_2(1) = 0.$$

**Guidelines:** We use the Lagrange form for the interpolation polynomial:

$$p_2(x) = \sum_{i=0}^2 y_i l_i(x), \quad l_i(x) := \prod_{j=0, j \neq i}^2 \frac{x - x_j}{x_i - x_j},$$

and with  $x_0 = 0$ ,  $x_1 = \frac{2}{3}$ ,  $x_2 = 1$ ,  $y_0 = y_2 = 0$ ,  $y_1 = 1$ .

## Problem 2

Construct the Lagrange interpolation polynomial  $p_1$  of degree 1 for a continuous function  $f$  defined on the interval  $[-1, 1]$ , using the interpolation points  $x_0 = -1$  and  $x_1 = 1$ . Show further that if the second order derivative of  $f$  exists and is continuous on  $[-1, 1]$  then

$$|f(x) - p_1(x)| \leq \frac{M_2}{2}(1 - x^2) \leq \frac{M_2}{2}, \quad x \in [-1, 1],$$

where

$$M_2 = \max_{x \in [-1, 1]} |f''(x)|.$$

Give an example of a function  $f$  for which equality is achieved.

**Guidelines:**

- Use the definition of Lagrange interpolation polynomial to find  $p_1$ .
- Use the error bound for the interpolation polynomial (Theorem 6.2).
- To find a concrete example for  $f$ , try with a quadratic polynomial of the form  $f(x) = ax^2 + bx + c$  and determine  $a$ ,  $b$  and  $c$ .

### Problem 3

Write down the Lagrange interpolation polynomial of degree 1 for the function  $f : x \mapsto x^3$ , using the points  $x_0 = 0$  and  $x_1 = a$ . Verify Theorem 6.2 by direct calculation showing that in this case  $\xi$  is unique and has the value  $\xi = \frac{1}{3}(x + a)$ .