

MA3203 - Exercise sheet 3

1. [1, Exercise III.8] Let $\Gamma = 1 \begin{array}{c} \xrightarrow{\alpha} \\ \xleftarrow{\beta} \end{array} 2$ be the Kronecker quiver. Define a representation H_λ of Γ by

$$H_\lambda = k \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{\lambda} \end{array} k .$$

for every $\lambda \in k$. Show that H_λ is indecomposable and that $H_\lambda \cong H_\mu$ if and only if $\lambda = \mu$. This shows that $k\Gamma$ is not of finite representation type when k is infinite.

2. [4, Exercise 2.1] Let $\Gamma = 1 \xrightarrow{\alpha} 2 \xleftarrow{\beta} 3$ and let

$$V = k^2 \begin{array}{c} \xrightarrow{\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}} \\ \xleftarrow{\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}} \end{array} k^2$$

. Write V as a direct sum of indecomposable representations of Γ .

3. Let $\Gamma = 1 \longrightarrow 2$ and let $V_1 \rightarrow V_2$ be a representation of Γ . Using Problem 2 b) in the previous exercise sheet, show that for any decomposition

$$V_1 \rightarrow V_2 \cong (k \rightarrow 0)^{m_1} \oplus (0 \rightarrow k)^{m_2} \oplus (k \xrightarrow{1} k)^{m_3}$$

of $V_1 \rightarrow V_2$ into indecomposable representations, the integers m_1, m_2, m_3 must be unique. This is a special case of the *Krull-Remak-Schmidt theorem*, which we will learn later in the course.

4. [2, Exercise 3.4] Let Γ be an arbitrary quiver and let V be a representation of Γ . Show that the following are equivalent.

- (a) V is indecomposable.

