

MA3203 - Exercise sheet 4

Throughout k denotes a field.

- [4, Problem 2.1] Find the representations corresponding to the modules Λe_i for the different possible values of i and for the different cases of Λ listed below. Also, find the representation corresponding to Λ (as a left Λ -module) in each case:

- (a) $\Lambda = k\Gamma/(\rho)$ where Γ is the quiver

$$1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3$$

and $\rho = \{\beta\alpha\}$.

- (b) $\Lambda = k\Gamma/(\rho)$ where Γ is the quiver

$$1 \xrightarrow{\alpha} 2 \begin{array}{c} \xrightarrow{\beta} \\ \xrightarrow{\gamma} \end{array} 3$$

and $\rho = \{\beta\alpha\}$.

- (c) $\Lambda = k\Gamma/(\rho)$ where Γ is the quiver

$$1 \begin{array}{c} \xrightarrow{\alpha} \\ \xrightarrow{\beta} \end{array} 2 \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \gamma$$

and $\rho = \{\gamma\alpha, \gamma^3\}$.

- Let $\Gamma = (\Gamma_0, \Gamma_1)$ be an arbitrary quiver. The two-sided ideal of $k\Gamma$ generated by Γ_1 is called the *arrow ideal* of $k\Gamma$, and is denoted R_Γ . An arbitrary two-sided ideal $I \subseteq k\Gamma$ is called *admissible* if there exists an integer $m \geq 2$ so that $R_\Gamma^m \subseteq I \subseteq R_\Gamma^2$.

- (a) Show that if I is admissible, then $k\Gamma/I$ is finite-dimensional (even if $k\Gamma$ is not).
- (b) [1, Exercise II.7a] Let $\Gamma = \alpha \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} 1 \begin{array}{c} \xrightarrow{\beta} \\ \xleftarrow{\gamma} \end{array} 2$ and let $\rho = \{\alpha^2, \gamma\beta, \beta\gamma - \beta\alpha\gamma\}$. Show that (ρ) is an admissible ideal, and compute $\dim_k k\Gamma/(\rho)$
- (c) [1, Example 2.7] Let $\Gamma = \alpha \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} 1 \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \beta$ and let $\rho = \{\beta\alpha, \beta^2\}$.

Show that (ρ) is not admissible and that $k\Gamma/(\rho)$ is infinite-dimensional. (This is an example of a ring which is left Noetherian but not right Noetherian due to J. Dieudonné, see [3, p16] for details. If you want a challenge you can try to prove this yourself).

3. ([2, Problem 4.1]) Let $\Gamma = \alpha \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} 1 \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \beta$ and let $\Lambda = k\Gamma/(\alpha^2, \beta^2, \beta\alpha)$.

- (a) Show that $(\alpha^2, \beta^2, \beta\alpha)$ is admissible, and compute the dimension of Λ .
- (b) Let $k\langle x, y \rangle$ be the ring of polynomials with coefficients in k and noncommuting variables x and y (that is, $xy \neq yx$). Show that $\Lambda \cong k\langle x, y \rangle / (x^2, y^2, yx)$.
- (c) Let $A = \left\{ \begin{bmatrix} a & 0 & 0 \\ b & a & 0 \\ c & d & a \end{bmatrix} : a, b, c, d \in k \right\}$. Show that A is a ring and that $\Lambda \cong A$.

4. Let Γ be the quiver $1 \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} \alpha$ and let $\rho = \{\alpha^n\}$. Using a similar argument as in the lecture, find all indecomposable representations of $k\Gamma/(\rho)$.

References

- [1] I. Assem, D. Simson, and A. Skowroński, *Elements of the Representation Theory of Associative Algebras 1: Techniques of Representation Theory*, London Math. Soc. Stud. Texts 65, Cambridge Univ. Press (2006).
- [2] E. Hanson, 2021 MA3203 Problem Sheets, NTNU, https://wiki.math.ntnu.no/ma3203/2021v/course_schedule.

- [3] H. Cartan, S. Eilenberg, *Homological Algebra*, Princeton University Press, Princeton, N. J., 1956
- [4] Ø. Solberg, 2017 MA3203 Problem Sheets, NTNU, <http://wiki.math.ntnu.no/ma3203/2017v/ovinge>.