

MA3203 - Exercise sheet 10

Throughout k denotes a field.

1. [1, Proposition 3.7 (a)-(c)] Let Λ be a ring and let L, M, N be Λ -modules. Show that the following hold:
 - (a) An element $m \in M$ belongs to $\text{rad } M$ if and only if $f(m) = 0$ for any $f \in \text{Hom}_\Lambda(M, S)$ and any simple Λ -module S .
 - (b) If $g: M \rightarrow N$ is a morphism of Λ -modules, then $g(\text{rad } M) \subseteq \text{rad } N$
 - (c) $\text{rad}(M \oplus N) = \text{rad}(M) \oplus \text{rad}(N)$.

2. [1, Exercise I.7.6] Let Λ be a ring, let M be a Λ -module, and let N be a Λ -submodule of M . Prove that
 - (a) $(N + \text{rad } M)/N \subseteq \text{rad}(M/N)$.
 - (b) If $N \subseteq \text{rad } M$, then $\text{rad}(M/N) = (\text{rad } M)/N$.

3. Find the radical of the module Λe_1 , where Λ is the path algebra of each of the following quivers (with relations):
 - (a) $\Lambda = k\Gamma$ for $\Gamma = 1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3$.
 - (b) $\Lambda = k\Gamma/(\gamma\alpha - \gamma\beta)$ for $\Gamma = 1 \begin{array}{c} \xrightarrow{\alpha} \\ \xrightarrow{\beta} \end{array} 2 \xrightarrow{\gamma} 3$.
 - (c) $\Lambda = k\Gamma$ for $\Gamma = 2 \xrightarrow{\alpha} 1 \xleftarrow{\beta} 3$.
 - (d) $\Lambda = k\Gamma/(\alpha\beta, \beta^2)$ for $\Gamma = 2 \xleftarrow{\alpha} 1 \begin{array}{c} \circlearrowleft \\ \beta \end{array}$.

