

**MA3204: HOMOLOGICAL ALGEBRA - EXERCISE SHEET 3**

**Exercise 1.** (The Yoneda embedding reflects exactness) Let  $\mathcal{A}$  be an abelian category and  $X \xrightarrow{f} Y \xrightarrow{g} Z$  a sequence in  $\mathcal{A}$ . Consider the following statements:

- (i) The sequence  $X \xrightarrow{f} Y \xrightarrow{g} Z$  is exact.
- (ii) For any object  $A$  in  $\mathcal{A}$ , the sequence

$$\text{Hom}_{\mathcal{A}}(A, X) \xrightarrow{f^*} \text{Hom}_{\mathcal{A}}(A, Y) \xrightarrow{g^*} \text{Hom}_{\mathcal{A}}(A, Z)$$

is exact.

- (iii) For any object  $A$  in  $\mathcal{A}$ , the sequence

$$\text{Hom}_{\mathcal{A}}(Z, A) \xrightarrow{g^*} \text{Hom}_{\mathcal{A}}(Y, A) \xrightarrow{f^*} \text{Hom}_{\mathcal{A}}(X, A)$$

is exact.

Show that (ii)  $\implies$  (i) and (iii)  $\implies$  (i).

**Exercise 2.** Let  $(F, G)$  be an adjoint pair of functors between the abelian categories  $\mathcal{A}$  and  $\mathcal{B}$ .

- (i) Show that  $F$  is right exact and  $G$  is left exact.
- (ii) Assume that the functor  $G$  is right exact. Show that the functor  $F$  preserves projective objects.
- (iii) Assume that the functor  $F$  is left exact. Show that the functor  $G$  preserves injective objects.

**Exercise 3.** Let  $\phi: R \rightarrow S$  be a homomorphism of rings. Show that  $(\phi^*, \phi_*, \phi^!)$ :

$$\begin{array}{ccc} & \phi^* & \\ & \curvearrowright & \\ \text{Mod-}S & \xrightarrow{\phi_*} & \text{Mod-}R \\ & \curvearrowleft & \\ & \phi^! & \end{array}$$

is an adjoint triple, i.e.  $(\phi^*, \phi_*)$  and  $(\phi_*, \phi^!)$  are adjoint pairs, where  $\phi_*$  is restriction of scalars,  $\phi^*(M) = M \otimes_R S$  is extension of scalars, and  $\phi^!(M) = \text{Hom}_R(S, M)$ .

**Exercise 4.** Let  $R$  be a ring and  $e^2 = e$  and idempotent element of  $R$ .

- (i) Show that the triple of functors  $(eR \otimes_{eRe} -, e(-), \text{Hom}_{eRe}(eR, -))$ :

$$\begin{array}{ccc} & eR \otimes_{eRe} - & \\ & \curvearrowright & \\ \text{Mod-}R & \xrightarrow{e(-)} & \text{Mod-}eRe \\ & \curvearrowleft & \\ & \text{Hom}_{eRe}(eR, -) & \end{array}$$

is an adjoint triple, where  $e(-)$  is multiplication with the idempotent element  $e$ .

- (ii) Show that the functor  $eR \otimes_{eRe} -: \text{Mod-}eRe \rightarrow \text{Mod-}R$  is fully faithful. Is the functor  $\text{Hom}_{eRe}(eR, -)$  fully faithful?
- (iii) What is the kernel of the functor  $e(-): \text{Mod-}R \rightarrow \text{Mod-}eRe$ ?

**Exercise 5.** Let  $R$  be a commutative ring and  $I, J$  two ideals in  $R$ . Show that  $R/I \otimes_R R/J \cong R/(I+J)$ . Using this, show that  $\mathbb{Z}_n \otimes_{\mathbb{Z}} \mathbb{Z}_m \cong \mathbb{Z}_d$  where  $d = \text{gcd}(m, n)$ .

**Exercise 6.** Let  $R$  and  $S$  be two  $k$ -algebras where  $k$  is a commutative ring. Show that  $R \otimes_k S$  is a  $k$ -algebra with multiplication:  $(r \otimes s) \cdot (r' \otimes s') = rr' \otimes ss'$ .

**Exercise 7.** Show that the  $k$ -algebras  $k[x, y]$  and  $k[x] \otimes_k k[y]$  are isomorphic.

**Exercise 8.** If  $R$  is a  $k$ -algebra, show that the  $k$ -algebras  $R \otimes_k M_n(k)$  and  $M_n(R)$  are isomorphic.

**Exercise 9.** Let  $R$  be a ring,  ${}_R A$  an  $R$ -module and  $(B_i)_{i \in I}$  a family of left  $R$ -modules.

(i) Show that we have the following isomorphisms:

$$\mathrm{Hom}_R(A, \prod_{i \in I} B_i) \cong \prod_{i \in I} \mathrm{Hom}_R(A, B_i)$$

and

$$\mathrm{Hom}_R(\prod_{i \in I} B_i, A) \cong \prod_{i \in I} \mathrm{Hom}_R(B_i, A)$$

(ii) Provide examples where the following groups are not isomorphic:

(a)  $\mathrm{Hom}_R(A, \prod_{i \in I} B_i) \not\cong \prod_{i \in I} \mathrm{Hom}_R(A, B_i)$ .

(b)  $\mathrm{Hom}_R(A, \prod_{i \in I} B_i) \not\cong \prod_{i \in I} \mathrm{Hom}_R(A, B_i)$ .

(c)  $\mathrm{Hom}_R(\prod_{i \in I} B_i, A) \not\cong \prod_{i \in I} \mathrm{Hom}_R(B_i, A)$ .

(d)  $\mathrm{Hom}_R(\prod_{i \in I} B_i, A) \not\cong \prod_{i \in I} \mathrm{Hom}_R(B_i, A)$ .

(iii) Show that if  ${}_R A$  is finitely generated, then there is an isomorphism:

$$\mathrm{Hom}_R(A, \prod_{i \in I} B_i) \cong \prod_{i \in I} \mathrm{Hom}_R(A, B_i)$$

**Exercise 10.** Exercises III.1 – III.4 from the notes.

CHRYSOSTOMOS PSAROUDAKIS, DEPARTMENT OF MATHEMATICAL SCIENCES, NORWEGIAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, 7491 TRONDHEIM, NORWAY

*E-mail address:* chrysostomos.psaroudakis@math.ntnu.no