TMA4115 Matematikk 3

Andrew Stacey

Norges Teknisk-Naturvitenskapelige Universitet
Trondheim

Spring 2010

Lecture 14: Determining Invertibility

Andrew Stacey

Norges Teknisk-Naturvitenskapelige Universitet
Trondheim

26th February 2010

Key Points

- More tests for invertibility
- Murky Waters

Recap

- Invertible matrix: A is invertible if $\exists B : AB = I_m$ and $BA = I_n$
- Learnt so far:
  - $A \longrightarrow B$ via Row Reduction:
    A invertible $\iff$ B invertible
  - $A \longrightarrow I_m$ via Row Reduction
    $\implies A$ invertible
  - Invertible $\implies$ Square

The Result of Row Reduction

Full Gauss–Jordan Row Reduction

Example outcomes:

The Result of Row Reduction

Full Gauss–Jordan Row Reduction

Example outcomes:

$$\begin{bmatrix} 1 & 2 & 0 & 0 & 4 \\ 0 & 0 & 1 & 0 & 6 \\ 0 & 0 & 0 & 1 & 8 \end{bmatrix}, \qquad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

The Result of Row Reduction

Full Gauss–Jordan Row Reduction

Example outcomes:

$$\begin{bmatrix} 1 & 2 & 0 & 0 & 4 \\ 0 & 0 & 1 & 0 & 6 \\ 0 & 0 & 0 & 1 & 8 \end{bmatrix}, \qquad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Neither is invertible

The Result of Row Reduction

Full Gauss–Jordan Row Reduction

Example outcomes:

$$\begin{bmatrix} 1 & 2 & 0 & 0 & 4 \\ 0 & 0 & 1 & 0 & 6 \\ 0 & 0 & 0 & 1 & 8 \end{bmatrix}, \qquad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

Neither is invertible

$$\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} \mapsto \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \qquad \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$$ impossible

Square Row Reduction Results

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & * \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Square Row Reduction Results

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & * \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

The first is the only matrix such that:

1. A is invertible

Square Row Reduction Results

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & * \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

The first is the only matrix such that:

1. A is invertible
   So A invertible $\iff$ row reduces to I

Square Row Reduction Results

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & * \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

The first is the only matrix such that:

1. A is invertible
   So A invertible $\iff$ row reduces to I
2. $A\mathbf{x} = \mathbf{b}$ has a solution for all $\mathbf{b}$

Square Row Reduction Results

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & * \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

The first is the only matrix such that:

1. A is invertible
   So A invertible $\iff$ row reduces to I
2. $A\mathbf{x} = \mathbf{b}$ has a solution for all $\mathbf{b}$
3. $A\mathbf{x} = \mathbf{b}$ has a unique solution for some $\mathbf{b}$

Square Row Reduction Results

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & * \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

The first is the only matrix such that:

1. A is invertible
   So A invertible $\iff$ row reduces to I
2. $A\mathbf{x} = \mathbf{b}$ has a solution for all $\mathbf{b}$
3. $A\mathbf{x} = \mathbf{b}$ has a unique solution for some $\mathbf{b}$
4. $A\mathbf{x} = \mathbf{0}$ has a unique solution

Square Row Reduction Results

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \begin{bmatrix} 1 & * \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

The first is the only matrix such that:

1. A is invertible
   So A invertible $\iff$ row reduces to I
2. $A\mathbf{x} = \mathbf{b}$ has a solution for all $\mathbf{b}$
3. $A\mathbf{x} = \mathbf{b}$ has a unique solution for some $\mathbf{b}$
4. $A\mathbf{x} = \mathbf{0}$ has a unique solution

This is independent of size of square matrix

Story So Far

A is invertible

$\Longleftrightarrow$

A is square and one (hence all) of:

1. A row reduces to I_m
2. $A\mathbf{x} = \mathbf{b}$ has a solution for all $\mathbf{b}$
3. $A\mathbf{x} = \mathbf{b}$ has a unique solution for some $\mathbf{b}$
4. $A\mathbf{x} = \mathbf{0}$ has a unique solution

Story So Far

A is invertible

$\Longleftrightarrow$

A is square and one (hence all) of:

1. A row reduces to I_m
2. $A\mathbf{x} = \mathbf{b}$ has a solution for all $\mathbf{b}$
3. $A\mathbf{x} = \mathbf{b}$ has a unique solution for some $\mathbf{b}$
4. $A\mathbf{x} = \mathbf{0}$ has a unique solution

Most important is . . .

Story So Far

A is invertible

$\Longleftrightarrow$

A is square and one (hence all) of:

1. A row reduces to I_m
2. $A\mathbf{x} = \mathbf{b}$ has a solution for all $\mathbf{b}$
3. $A\mathbf{x} = \mathbf{b}$ has a unique solution for some $\mathbf{b}$
4. $A\mathbf{x} = \mathbf{0}$ has a unique solution

Most important is . . .

Computing The Inverse

The Amazing Gauss

GE is so amazing you can even find the inverse using it.

Computing The Inverse

The Amazing Gauss

GE is so amazing you can even find the inverse using it.

Idea

- A invertible

Computing The Inverse

The Amazing Gauss

GE is so amazing you can even find the inverse using it.

Idea

- A invertible $\iff$ A row reduces to I_m

Computing The Inverse

The Amazing Gauss

GE is so amazing you can even find the inverse using it.

Idea

- A invertible $\iff$ A row reduces to I_m
- Row ops $\longleftrightarrow$ elementary matricces E_j:

$$E_k \cdots E_1 A = I_m$$

Computing The Inverse

The Amazing Gauss

GE is so amazing you can even find the inverse using it.

Idea

- A invertible $\iff$ A row reduces to I_m
- Row ops $\longleftrightarrow$ elementary matricces E_j:

$$E_k \cdots E_1 A = I_m$$

- Apply same row ops to I_m to get A^{-1}!

$$E_k \cdots E_1 I_m = E_k \cdots E_1 A A^{-1} = A^{-1}$$

Computing The Inverse

The Amazing Gauss

GE is so amazing you can even find the inverse using it.

Idea

- A invertible $\iff$ A row reduces to I_m
- Row ops $\longleftrightarrow$ elementary matricces E_j:

$$E_k \cdots E_1 A = I_m$$

- Apply same row ops to I_m to get A^{-1}!

$$E_k \cdots E_1 I_m = E_k \cdots E_1 A A^{-1} = A^{-1}$$

- In practice: apply row ops to $[A|I_m]$.

Computing The Inverse

The Amazing Gauss

GE is so amazing you can even find the inverse using it.

Idea

- A invertible $\iff$ A row reduces to I_m
- Row ops $\longleftrightarrow$ elementary matricces E_j:

$$E_k \cdots E_1 A = I_m$$

- Apply same row ops to I_m to get A^{-1}!

$$E_k \cdots E_1 I_m = E_k \cdots E_1 A A^{-1} = A^{-1}$$

- In practice: apply row ops to $[A|I_m]$.
  Result is $[I_m|B] \iff$ A invertible and $A^{-1} = B$

Example

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

Example

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 1 & 3 & 3 & 0 & 1 & 0 \\ 1 & 4 & 6 & 0 & 0 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 1 & 3 & 3 & 0 & 1 & 0 \\ 1 & 4 & 6 & 0 & 0 & 1 \end{array}\right] \begin{array}{c} R_2 - R_1 \\ \longrightarrow \\ R_3 - R_1 \end{array} \left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 2 & 5 & -1 & 0 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 1 & 3 & 3 & 0 & 1 & 0 \\ 1 & 4 & 6 & 0 & 0 & 1 \end{array}\right] \xrightarrow[R_3 - R_1]{R_2 - R_1} \left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 2 & 5 & -1 & 0 & 1 \end{array}\right]$$

$$\xrightarrow[R_3 - 2R_2]{R_1 - 2R_2} \left[\begin{array}{ccc|ccc} 1 & 0 & -3 & 3 & -2 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 1 & 3 & 3 & 0 & 1 & 0 \\ 1 & 4 & 6 & 0 & 0 & 1 \end{array}\right] \begin{array}{c} R_2 - R_1 \\ \longrightarrow \\ R_3 - R_1 \end{array} \left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 2 & 5 & -1 & 0 & 1 \end{array}\right]$$

$$\begin{array}{c} R_1 - 2R_2 \\ \longrightarrow \\ R_3 - 2R_2 \end{array} \left[\begin{array}{ccc|ccc} 1 & 0 & -3 & 3 & -2 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{array}\right]$$

$$\begin{array}{c} R_1 + 3R_3 \\ \longrightarrow \\ R_2 - 2R_3 \end{array} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 6 & -8 & 3 \\ 0 & 1 & 0 & -3 & 5 & -2 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 3 & 3 \\ 1 & 4 & 6 \end{bmatrix} \qquad \text{Yes, } A^{-1} = \begin{bmatrix} 6 & -8 & 3 \\ -3 & 5 & -2 \\ 1 & -2 & 1 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 1 & 3 & 3 & 0 & 1 & 0 \\ 1 & 4 & 6 & 0 & 0 & 1 \end{array}\right] \begin{array}{c} R_2 - R_1 \\ \longrightarrow \\ R_3 - R_1 \end{array} \left[\begin{array}{ccc|ccc} 1 & 2 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 2 & 5 & -1 & 0 & 1 \end{array}\right]$$

$$\begin{array}{c} R_1 - 2R_2 \\ \longrightarrow \\ R_3 - 2R_2 \end{array} \left[\begin{array}{ccc|ccc} 1 & 0 & -3 & 3 & -2 & 0 \\ 0 & 1 & 2 & -1 & 1 & 0 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{array}\right]$$

$$\begin{array}{c} R_1 + 3R_3 \\ \longrightarrow \\ R_2 - 2R_3 \end{array} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 6 & -8 & 3 \\ 0 & 1 & 0 & -3 & 5 & -2 \\ 0 & 0 & 1 & 1 & -2 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

Example

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array}\right] \begin{array}{c} R_2 - 4R_1 \\ \longrightarrow \\ R_3 - 7R_1 \end{array} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array}\right] \xrightarrow[R_3 - 7R_1]{R_2 - 4R_1} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array}\right]$$

$$\xrightarrow{R_3 - 2R_2} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array}\right] \xrightarrow[R_3 - 7R_1]{R_2 - 4R_1} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array}\right]$$

$$\xrightarrow{R_3 - 2R_2} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array}\right]$$

$$\longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 0 & -2\tfrac{2}{3} & 1\tfrac{2}{3} \\ 0 & 1 & 2 & 0 & 2\tfrac{1}{3} & -1\tfrac{1}{3} \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array}\right]$$

Example

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} \qquad \text{No, } A \begin{bmatrix} 1 \\ -2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Is A invertible, if so what is A^{-1}?

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 4 & 5 & 6 & 0 & 1 & 0 \\ 7 & 8 & 9 & 0 & 0 & 1 \end{array}\right] \xrightarrow[R_3 - 7R_1]{R_2 - 4R_1} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & -6 & -12 & -7 & 0 & 1 \end{array}\right]$$

$$\xrightarrow{R_3 - 2R_2} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & -3 & -6 & -4 & 1 & 0 \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array}\right]$$

$$\longrightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 0 & -2\tfrac{2}{3} & 1\tfrac{2}{3} \\ 0 & 1 & 2 & 0 & 2\tfrac{1}{3} & -1\tfrac{1}{3} \\ 0 & 0 & 0 & 1 & -2 & 1 \end{array}\right]$$

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?
Can solve $A\mathbf{x} = \mathbf{b}$ for all $\mathbf{b}$

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?
Can solve $A\mathbf{x} = \mathbf{b}$ for all $\mathbf{b}$
Apply GE ($a \neq 0$):

$$\left[\begin{array}{cc|c} a & b & b_1 \\ c & d & b_2 \end{array}\right]$$

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?
Can solve $A\mathbf{x} = \mathbf{b}$ for all $\mathbf{b}$
Apply GE ($a \neq 0$):

$$\left[\begin{array}{cc|c} a & b & b_1 \\ c & d & b_2 \end{array}\right] \xrightarrow[a \neq 0]{R_2 - {}^{c}/_{a} R_1} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & d - {}^{cb}/_{a} & b_2 - {}^{c}/_{a} b_1 \end{array}\right]$$

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?
Can solve $A\mathbf{x} = \mathbf{b}$ for all $\mathbf{b}$
Apply GE ($a \neq 0$):

$$\left[\begin{array}{cc|c} a & b & b_1 \\ c & d & b_2 \end{array}\right] \xrightarrow[a \neq 0]{R_2 - c/a R_1} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & d - cb/a & b_2 - c/a b_1 \end{array}\right]$$

$$\xrightarrow{R_2 \to aR_2} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & ad - cb & ab_2 - cb_1 \end{array}\right]$$

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?
Can solve $A\mathbf{x} = \mathbf{b}$ for all $\mathbf{b}$
Apply GE ($a \neq 0$):

$$\left[\begin{array}{cc|c} a & b & b_1 \\ c & d & b_2 \end{array}\right] \xrightarrow[a \neq 0]{R_2 - c/a R_1} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & d - cb/a & b_2 - c/a b_1 \end{array}\right]$$

$$\xrightarrow{R_2 \to aR_2} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & ad - cb & ab_2 - cb_1 \end{array}\right]$$

If $a \neq 0$ then A invertible $\iff$ $ad - bc \neq 0$

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?
Can solve $A\mathbf{x} = \mathbf{b}$ for all $\mathbf{b}$
Apply GE ($a \neq 0$):

$$\left[\begin{array}{cc|c} a & b & b_1 \\ c & d & b_2 \end{array}\right] \xrightarrow[a \neq 0]{R_2 - c/a R_1} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & d - cb/a & b_2 - c/a b_1 \end{array}\right]$$

$$\xrightarrow{R_2 \to aR_2} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & ad - cb & ab_2 - cb_1 \end{array}\right]$$

If $a \neq 0$ then A invertible $\iff$ $ad - bc \neq 0$
If $c \neq 0$ then swap so A invertible $\iff$ $cb - ad \neq 0$

Other Methods

When is

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

invertible?
Can solve $A\mathbf{x} = \mathbf{b}$ for all $\mathbf{b}$
Apply GE ($a \neq 0$):

$$\left[\begin{array}{cc|c} a & b & b_1 \\ c & d & b_2 \end{array}\right] \xrightarrow[a\neq 0]{R_2 - c/a R_1} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & d - cb/a & b_2 - c/a b_1 \end{array}\right]$$

$$\xrightarrow{R_2 \to aR_2} \left[\begin{array}{cc|c} a & b & b_1 \\ 0 & ad - cb & ab_2 - cb_1 \end{array}\right]$$

If $a \neq 0$ then A invertible $\iff$ $ad - bc \neq 0$
If $c \neq 0$ then swap so A invertible $\iff$ $cb - ad \neq 0$
If $a = c = 0$ then A not invertible and $cb - ad = 0$

It's that Strange Sense of Deja Vu

y_1, y_2 solutions of an ODE
linearly independent

$$\Longleftrightarrow$$

$$y_1 y_2' - y_1' y_2 \neq 0$$

A
invertible

$$\Longleftrightarrow$$

$$ad - bc \neq 0$$

It's that Strange Sense of Deja Vu

y_1, y_2 solutions of an ODE
linearly independent

$$\Longleftrightarrow$$

$$y_1 y_2' - y_1' y_2 \neq 0$$

$$A$$

invertible

$$\Longleftrightarrow$$

$$ad - bc \neq 0$$

Not a coincidence!

Determining Invertibility

Definition

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

the determinant of A is

$$\det(A) := \begin{vmatrix} a & b \\ c & d \end{vmatrix} := ad - bc$$

Determining Invertibility

Definition

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

the determinant of A *is*

$$\det(A) := \begin{vmatrix} a & b \\ c & d \end{vmatrix} := ad - bc$$

Lemma

$$A \text{ invertible} \iff \det(A) \neq 0$$

Bigger and . . . Better?

Sort of Definition

Square matrices of any size have determinants.

Bigger and . . . Better?

Sort of Definition

Square matrices of any size have determinants.

Exact definition not exactly . . . memorable:

$$\det(A) = \sum_{\sigma \in \Sigma_n} (-1)^{\sigma} a_{11\sigma} \cdots a_{nn\sigma}$$

Bigger and . . . Better?

Sort of Definition

Square matrices of any size have determinants.

Exact definition not exactly . . . memorable:

$$\det(A) = \sum_{\sigma \in \Sigma_n} (-1)^{\sigma} a_{11\sigma} \cdots a_{nn\sigma}$$

Best: Learn rules for computation.

Don't Know? Ask Gauss

Key Properties for Computation

1. $\det(I_n) = 1$

Don't Know? Ask Gauss

Key Properties for Computation

1. $\det(I_n) = 1$
2. $\det(AB) = \det(A)\det(B)$

Don't Know? Ask Gauss

Key Properties for Computation

1. $\det(I_n) = 1$
2. $\det(AB) = \det(A)\det(B)$
3. Triangular matrices:

$$\begin{vmatrix} 2 & 3 & 4 \\ 0 & 5 & 6 \\ 0 & 0 & 7 \end{vmatrix} = 2 \times 5 \times 7 = 70$$

Don't Know? Ask Gauss

Key Properties for Computation

1. $\det(I_n) = 1$
2. $\det(AB) = \det(A)\det(B)$
3. Triangular matrices:

$$\begin{vmatrix} 2 & 3 & 4 \\ 0 & 5 & 6 \\ 0 & 0 & 7 \end{vmatrix} = 2 \times 5 \times 7 = 70$$

4. Row Swaps:

$$\begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = - \begin{vmatrix} 4 & 5 \\ 2 & 3 \end{vmatrix}$$

Don't Know? Ask Gauss

Key Properties for Computation

1. $\det(I_n) = 1$
2. $\det(AB) = \det(A)\det(B)$
3. Triangular matrices:

$$\begin{vmatrix} 2 & 3 & 4 \\ 0 & 5 & 6 \\ 0 & 0 & 7 \end{vmatrix} = 2 \times 5 \times 7 = 70$$

4. Row Swaps:

$$\begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = - \begin{vmatrix} 4 & 5 \\ 2 & 3 \end{vmatrix}$$

5. Row Multipliers:

$$\begin{vmatrix} 2 & 4 \\ 3 & 5 \end{vmatrix} = 2 \begin{vmatrix} 1 & 2 \\ 3 & 5 \end{vmatrix}$$

Note: only one row affected.

Don't Know? Ask Gauss

6. Row Adds:

$$\begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} 2 & 3 \\ 0 & -1 \end{vmatrix}$$

No change!

Don't Know? Ask Gauss

6. Row Adds:

$$\begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} 2 & 3 \\ 0 & -1 \end{vmatrix}$$

   No change!

7. Works for columns too.

Don't Know? Ask Gauss

6. Row Adds:

$$\begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} 2 & 3 \\ 0 & -1 \end{vmatrix}$$

   No change!

7. Works for columns too.

Conclusion

Whenever you need to compute something, use GE!

Don't Know? Ask Gauss

6. Row Adds:

$$\begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} 2 & 3 \\ 0 & -1 \end{vmatrix}$$

   No change!

7. Works for columns too.

Conclusion

Whenever you need to compute something, use GE!

Mnemonic

$$\begin{vmatrix} 0 & 1 \\ 2 & 2 \end{vmatrix} = -2$$

Example

$$\begin{vmatrix} 0 & 2 & 3 \\ 3 & 4 & 0 \\ 1 & 6 & 2 \end{vmatrix}$$

Example

$$\begin{vmatrix} 0 & 2 & 3 \\ 3 & 4 & 0 \\ 1 & 6 & 2 \end{vmatrix} = - \begin{vmatrix} 1 & 6 & 2 \\ 3 & 4 & 0 \\ 0 & 2 & 3 \end{vmatrix} (R_1 \leftrightarrow R_3)$$

Example

$$\begin{vmatrix} 0 & 2 & 3 \\ 3 & 4 & 0 \\ 1 & 6 & 2 \end{vmatrix} = -\begin{vmatrix} 1 & 6 & 2 \\ 3 & 4 & 0 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_1 \leftrightarrow R_3)$$

$$= -\begin{vmatrix} 1 & 6 & 2 \\ 0 & -14 & -6 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_2 - 3R_1)$$

Example

$$\begin{vmatrix} 0 & 2 & 3 \\ 3 & 4 & 0 \\ 1 & 6 & 2 \end{vmatrix} = -\begin{vmatrix} 1 & 6 & 2 \\ 3 & 4 & 0 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_1 \leftrightarrow R_3)$$

$$= -\begin{vmatrix} 1 & 6 & 2 \\ 0 & -14 & -6 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_2 - 3R_1) \qquad = -2\begin{vmatrix} 1 & 3 & 2 \\ 0 & -7 & -6 \\ 0 & 1 & 3 \end{vmatrix} \quad (C_2 \to {}^1\!/_2 C_2)$$

Example

$$\begin{vmatrix} 0 & 2 & 3 \\ 3 & 4 & 0 \\ 1 & 6 & 2 \end{vmatrix} = -\begin{vmatrix} 1 & 6 & 2 \\ 3 & 4 & 0 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_1 \leftrightarrow R_3)$$

$$= -\begin{vmatrix} 1 & 6 & 2 \\ 0 & -14 & -6 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_2 - 3R_1) \qquad = -2\begin{vmatrix} 1 & 3 & 2 \\ 0 & -7 & -6 \\ 0 & 1 & 3 \end{vmatrix} \quad (C_2 \to {}^1\!/_2 C_2)$$

$$= 2\begin{vmatrix} 1 & 3 & 2 \\ 0 & 1 & 3 \\ 0 & -7 & -6 \end{vmatrix} \quad (R_2 \leftrightarrow R_3)$$

Example

$$\begin{vmatrix} 0 & 2 & 3 \\ 3 & 4 & 0 \\ 1 & 6 & 2 \end{vmatrix} = -\begin{vmatrix} 1 & 6 & 2 \\ 3 & 4 & 0 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_1 \leftrightarrow R_3)$$

$$= -\begin{vmatrix} 1 & 6 & 2 \\ 0 & -14 & -6 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_2 - 3R_1) \qquad = -2\begin{vmatrix} 1 & 3 & 2 \\ 0 & -7 & -6 \\ 0 & 1 & 3 \end{vmatrix} \quad (C_2 \to {}^1\!/_2 C_2)$$

$$= 2\begin{vmatrix} 1 & 3 & 2 \\ 0 & 1 & 3 \\ 0 & -7 & -6 \end{vmatrix} \quad (R_2 \leftrightarrow R_3) \qquad = 2\begin{vmatrix} 1 & 3 & 2 \\ 0 & 1 & 3 \\ 0 & 0 & 15 \end{vmatrix} \quad (R_3 + 7R_2)$$

Example

$$\begin{vmatrix} 0 & 2 & 3 \\ 3 & 4 & 0 \\ 1 & 6 & 2 \end{vmatrix} = -\begin{vmatrix} 1 & 6 & 2 \\ 3 & 4 & 0 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_1 \leftrightarrow R_3)$$

$$= -\begin{vmatrix} 1 & 6 & 2 \\ 0 & -14 & -6 \\ 0 & 2 & 3 \end{vmatrix} \quad (R_2 - 3R_1) \qquad = -2\begin{vmatrix} 1 & 3 & 2 \\ 0 & -7 & -6 \\ 0 & 1 & 3 \end{vmatrix} \quad (C_2 \to {}^1\!/_2 C_2)$$

$$= 2\begin{vmatrix} 1 & 3 & 2 \\ 0 & 1 & 3 \\ 0 & -7 & -6 \end{vmatrix} \quad (R_2 \leftrightarrow R_3) \qquad = 2\begin{vmatrix} 1 & 3 & 2 \\ 0 & 1 & 3 \\ 0 & 0 & 15 \end{vmatrix} \quad (R_3 + 7R_2)$$

$$= 2 \times 15 = 30$$

Story So Far

- To compute anything:
  Use Gaussian Elimination

Muddiest Point

The Winner of the Muddiest Point Competition is

Muddiest Point

The Winner of the Muddiest Point Competition is

Inhomogeneous ODEs

Recall

$$y'' + p(x)y' + q(x)y = r(x)$$

Recall

$$y'' + p(x)y' + q(x)y = r(x)$$

Examples

1. $y'' + y' - 2y = \sin(x)$

Recall

$$y'' + p(x)y' + q(x)y = r(x)$$

Examples

1. $y'' + y' - 2y = \sin(x)$
2. $x^2y'' + xy' - y = x^2 + x^3$

Recall

$$y'' + p(x)y' + q(x)y = r(x)$$

Examples

1. $y'' + y' - 2y = \sin(x)$
2. $x^2y'' + xy' - y = x^2 + x^3$
3. $y'' + x^{-2}y' - x^{-3}y = x^{-3}$

General Method

1. Solve homogeneous ODE, y_1, y_2

General Method

1. Solve homogeneous ODE, y_1, y_2

2. Find a solution of inhomogeneous ODE, y_p

General Method

1. Solve homogeneous ODE, y_1, y_2

2. Find a solution of inhomogeneous ODE, y_p

3. Fit to initial values:

$$y_p(x_0) + a y_1(x_0) + b y_2(x_0)$$
$$y_p'(x_0) + a y_1'(x_0) + b y_2'(x_0)$$

General Method

1. Solve homogeneous ODE, y_1, y_2
   1.1 "Well-known ODE" $\Longrightarrow$ "look up" solution

2. Find a solution of inhomogeneous ODE, y_p

3. Fit to initial values:

$$y_p(x_0) + a y_1(x_0) + b y_2(x_0)$$
$$y_p'(x_0) + a y_1'(x_0) + b y_2'(x_0)$$

General Method

1. Solve homogeneous ODE, y_1, y_2
   1.1 "Well-known ODE" $\Longrightarrow$ "look up" solution
   1.2 Otherwise, guess one solution

2. Find a solution of inhomogeneous ODE, y_p

3. Fit to initial values:

$$y_p(x_0) + a y_1(x_0) + b y_2(x_0)$$
$$y_p'(x_0) + a y_1'(x_0) + b y_2'(x_0)$$

General Method

1. Solve homogeneous ODE, y_1, y_2
   - 1.1 "Well-known ODE" $\Longrightarrow$ "look up" solution
   - 1.2 Otherwise, guess one solution
   - 1.3 "Reduction of order" for other
2. Find a solution of inhomogeneous ODE, y_p

3. Fit to initial values:

$$y_p(x_0) + a y_1(x_0) + b y_2(x_0)$$
$$y_p'(x_0) + a y_1'(x_0) + b y_2'(x_0)$$

General Method

1. Solve homogeneous ODE, y_1, y_2
   - 1.1 "Well-known ODE" $\Longrightarrow$ "look up" solution
   - 1.2 Otherwise, guess one solution
   - 1.3 "Reduction of order" for other
2. Find a solution of inhomogeneous ODE, y_p
   - 2.1 Guess

3. Fit to initial values:

$$y_p(x_0) + a y_1(x_0) + b y_2(x_0)$$
$$y_p'(x_0) + a y_1'(x_0) + b y_2'(x_0)$$

General Method

1. Solve homogeneous ODE, y_1, y_2
   - 1.1 "Well-known ODE" $\Longrightarrow$ "look up" solution
   - 1.2 Otherwise, guess one solution
   - 1.3 "Reduction of order" for other
2. Find a solution of inhomogeneous ODE, y_p
   - 2.1 Guess
   - 2.2 Otherwise, use "variation of parameters"
3. Fit to initial values:

$$y_p(x_0) + a y_1(x_0) + b y_2(x_0)$$
$$y_p'(x_0) + a y_1'(x_0) + b y_2'(x_0)$$

Examples

$$y'' + y' - 2y = \sin(x)$$

Examples

$$x^2y'' + xy' - y = x^2 + x^3$$

Examples

$$y'' + x^{-2}y' - x^{-3}y = x^{-3}$$

Summary

- Matrices:
  Use GE whenever you want an actual answer
- ODEs:
  Use intelligent guesswork