

NTNU
Norwegian University of
Science and Technology

TMA4115 - Calculus 3

Week 15, part 1

Louise Meier Carlsen
Norwegian University of Science and Technology

Spring 2013

Wednesday's lecture

Yesterday we introduced and studied

- *eigenvectors*, *eigenvalues* and *eigenspaces* of square matrices (Section 5.1),
- the *characteristic polynomial* of a square matrix (Section 5.2).

Today's lecture

Today we shall look at

- *diagonal* matrices,
- how to *diagonalizable* a matrix.

Eigenvectors, eigenvalues and eigenspaces

Let A be an $n \times n$ matrix. Recall that

- An *eigenvalue* of A is a scalar λ such that the equation $A\mathbf{x} = \lambda\mathbf{x}$ has a nontrivial solution.
- If λ is an eigenvalue of A, then a nontrivial solution to the equation $A\mathbf{x} = \lambda\mathbf{x}$ is called an *eigenvector (of A) corresponding to* λ.
- If λ is an eigenvalue of A, then the set of solutions to the equation $A\mathbf{x} = \lambda\mathbf{x}$ is called the *eigenspace (of A) corresponding to* λ.

NTNU
Norwegian University of
Science and Technology

Finding a basis of an eigenspace

Let A be an $n \times n$ matrix and let λ be an eigenvalue of A. Recall that the eigenspace of A corresponding to λ is equal to $\text{Nul}(A - \lambda I_n)$, so we can find a basis for the eigenspace of A corresponding to λ by:

1. row reducing $A - \lambda I_n$ to its reduced echelon form,
2. write the solutions to the equation $(A - \lambda I_n)\mathbf{x} = \mathbf{0}$ (which is equivalent to the equation $A\mathbf{x} - \mathbf{0}$) as a linear combinations of vectors using the free variables as parameters.
3. Then these vectors form a basis of the eigenspace of A corresponding to λ.

NTNU
Norwegian University of
Science and Technology

Finding the eigenvalues of a matrix

Let A be an $n \times n$ matrix. Recall that a scalar λ is an eigenvalue of A if and only if it is a zero of the *characteristic polynomial* $\det(A - \lambda I_n)$ of A.

Example

Let us find the eigenvalues of the matrix $A = \begin{bmatrix} 1 & -3 & 6 & 0 \\ 0 & -2 & 1 & 1 \\ 0 & 0 & 0 & 2 \\ 0 & 0 & 0 & 7 \end{bmatrix}$.

The eigenvalues of a triangular matrix

Theorem 1

The eigenvalues of a triangular matrix are the entries on its main diagonal.

Example

Let $A = \begin{bmatrix} 1 & 2 & -1 \\ -5 & 6 & -3 \\ 5 & 2 & 11 \end{bmatrix}$, $D = \begin{bmatrix} 4 & -8 & 4 \\ 0 & 6 & 4 \\ 0 & 0 & 8 \end{bmatrix}$ and

$P = \begin{bmatrix} 1 & 1 & -1 \\ 1 & -1 & 1 \\ -1 & 1 & 1 \end{bmatrix}$.

NTNU
Norwegian University of
Science and Technology

Similarity

Let A and B be $n \times n$ matrices.

- A is *similar* to B if there is an invertible matrix P such that $P^{-1}AP = B$.
- If A is similar to B, then B is similar to A (because if $P^{-1}AP = B$, and we let $Q = P^{-1}$, then $Q^{-1}BQ = PBP^{-1} = PP^{-1}BP^{-1}P = A$).
- If A is similar to B (and thus B is similar to A), then we say that A and B *are similar*.

NTNU
Norwegian University of
Science and Technology

The characteristic polynomials of similar matrices

Theorem 4

If two matrices A and B are similar, then they have the same characteristic polynomial and hence the same eigenvalues (with the same multiplicity).

The *multiplicity* of an eigenvalue λ_k of an $n \times n$ matrix A is the maximal m such that $(\lambda - \lambda_k)^m$ is a factor of $\det(A - \lambda I_n)$.

NTNU
Norwegian University of
Science and Technology

Warnings

- Matrices that have the same eigenvalues are not necessarily similar.
- Row operations on a matrix usually changes its eigenvalues.

NTNU
Norwegian University of
Science and Technology

Finding eigenvalues via the QR algorithm

The previous theorem can be used to calculate the eigenvalues of a square matrix.

- If A is an $n \times n$-matrix, then there exists under certain conditions a sequence $A_1, A_2, \ldots$ of $n \times n$-matrix such that each A_k is similar to A, and such that A_k converges to a triangular matrix B.
- Since each A_k is similar to A, the eigenvalues of each A_k, and hence of B, are the same as the eigenvalues of A.
- So the diagonal entries of A_k converge to the eigenvalues of A.

NTNU
Norwegian University of
Science and Technology

Diagonalizable matrices

- A *diagonal* matrix is a square matrix whose entries outside the main diagonal are all zero.
- If D is a diagonal matrix, then the eigenvalues of D are the entries on the main diagonal.
- If D is a diagonal matrix, then $\mathbf{e}_i$ is an eigenvector of D corresponding to the i'th entry on the main diagonal.
- A matrix A is *diagonalizable* if it is similar to a diagonal matrix (that is, if $A = PDP^{-1}$ for some invertible matrix P and some diagonal matrix D).

NTNU
Norwegian University of
Science and Technology

The diagonalization Theorem

Theorem 5

An $n \times n$ matrix A is diagonalizable if and only if A has n linearly independent eigenvectors.
In fact, $A = PDP^{-1}$, with D a diagonal matrix, if and only if the columns of P are n linearly independent eigenvectors of A, in which case the diagonal entries of D are eigenvalues of A corresponding, respectively, to the eigenvectors in P.

NTNU
Norwegian University of
Science and Technology

How to diagonalize a matrix (if it is diagonalizable)

1. Find the eigenvalues of A.
2. Find n linearly independent eigenvectors of A (if possible).
3. Construct P from the linearly independent eigenvectors of A.
4. Construct D from the corresponding eigenvalues.
5. Then $A = PDP^{-1}$.

NTNU
Norwegian University of
Science and Technology

Example

Let us determine if $A = \begin{bmatrix} 0 & 2 \\ -3 & 5 \end{bmatrix}$ is diagonalizable and diagonalize it if it is and find an expression for A^n.

Matrices with n distinct eigenvalues are diagonalizable

Theorem 6

An $n \times n$ matrix with n distinct eigenvalues is diagonalizable.

NTNU
Norwegian University of
Science and Technology

Matrices with fewer than n distinct eigenvalues

Theorem 7

Let A be an $n \times n$ matrix whose distinct eigenvalues are $\lambda_1, \ldots, \lambda_p$.

1. For $1 \leq k \leq p$, the dimension of the eigenspace for λ_k is less than or equal to the multiplicity of the eigenvalue λ_k.

NTNU
Norwegian University of
Science and Technology

Matrices with fewer than n distinct eigenvalues

Theorem 7

2. A is diagonalizable if and only if the sum of the dimensions of the eigenspaces equals n, and this happens if and only if the characteristic polynomial factors completely into linear factors and the dimension of the eigenspace for each λ_k equals the multiplicity of λ_k.
3. If A is diagonalizable and $\mathcal{B}_k$ is a basis for the eigenspace corresponding to λ_k for each k, then the total collection of vectors in $\mathcal{B}_1, \ldots, \mathcal{B}_k$ forms an eigenvector basis for $\mathbb{R}^n$.

NTNU
Norwegian University of Science and Technology

Problem 7 from the exam from June 2012

1. Find all the eigenvalues of $A = \begin{bmatrix} 2 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & -1 & 4 \end{bmatrix}$.
2. Find a basis for each eigenspace of A. Is A diagonalisable?

NTNU
Norwegian University of
Science and Technology

Problem 11 from August 2004

Let A be a diagonalizable matrix where every eigenvalue is either 1 or -1. Show that $A^{-1} = A$.

NTNU
Norwegian University of
Science and Technology

Problem 7 from December 2005

Let A and B be two $n \times n$-matrices.

1. Show that if 0 is an eigenvalue of AB, then 0 is an eigenvalue of BA.
2. Show that if B is invertible and λ is an eigenvalue of AB, then λ is an eigenvalue of BA.

NTNU
Norwegian University of
Science and Technology

Review quiz

Let A be a square matrix. Decide for each of the following statements, whether the statement is true or false.

1. The sum of two eigenvectors of a matrix A is also an eigenvector of A.
2. Two eigenvectors corresponding to the same eigenvalue are always linearly dependent.
3. If A and B are invertible $n \times n$ matrices, then AB is similar to BA.
4. Each eigenvector of A is also an eigenvector of A^2.
5. If A is an $n \times n$ diagonalizable matrix, then each vector in $\mathbb{R}^n$ can be written as a linear combination of eigenvectors of A.
6. If A contains a row or column of zeros, then 0 is an eigenvalue of A.

NTNU
Norwegian University of
Science and Technology

Plan for Wednesday

On Wednesday we shall study

- real matrices with complex eigenvalues,
- systems of first order differential equations.

Sections 5.5 and 5.7 in “Linear Algebras and Its Applications” (pages 295-301, and 311-319), and Section 4.2 in “Second-Order Equations” (pages xlv-xlix).

NTNU
Norwegian University of
Science and Technology