

NTNU
Norwegian University of
Science and Technology

TMA4115 - Calculus 3

Lecture 1, Jan 16

Toke Meier Carlsen
Norwegian University of Science and Technology

Spring 2013

Course web page

Information about the course can be found at
`http://wiki.math.ntnu.no/tma4115/2013v`.

NTNU
Norwegian University of
Science and Technology

Overview of the course

The topics of the course are

- Complex numbers (1.5 weeks)
- Second order linear differential equations (1.5 weeks)
- Linear algebra (10 weeks)

NTNU
Norwegian University of
Science and Technology

Complex numbers

- A complex number is a number which can be written as $a + ib$ where a and b are real numbers and i satisfies $i^2 = -1$.
- The Italian mathematician Gerolamo Cardano is credited with introducing complex numbers in around 1545 in order to solve cubic equations.
- Complex numbers are not only used for solving polynomial equations, but naturally show up all over in mathematics and are used in many scientific fields, including engineering, electromagnetism and quantum physics.

NTNU
Norwegian University of
Science and Technology

Complex numbers as points in the plane

- A complex number can be represented as a point (a, b) in the plane.
- If $z = (a, b)$, then a is called the *real part* of z and is denoted by $\mathrm{Re}(z)$.
- b is called the *imaginary part* of z and is denoted by $\mathrm{Im}(z)$.
- The length $\sqrt{a^2 + b^2}$ of the line from $(0, 0)$ to (a, b) is called the *modulus* or the *absolute value* of z and is denoted by $|z|$.



NTNU
Norwegian University of Science and Technology

Complex numbers as points in the plane

- The angle between the line through $(0,0)$ and (a,b) and the positive part of the real axis is called the *argument* of z and is denoted by $\arg(z)$.
- $\arg(z)$ is not unique. If $\theta = \arg(z)$, then also $\theta + 2\pi = \arg(z)$.



If we want to be precise, then $\arg(z)$ is really the set of all angles θ which satisfies that if we rotate the positive part of the real axis by θ, then it lands on the line through $(0,0)$ and (a,b).

NTNU
Norwegian University of Science and Technology

Complex numbers as points in the plane

- The unique value of $\arg(z)$ in the interval $(-\pi, \pi]$ is called the *principal argument* of z and is denoted by $\mathrm{Arg}(z)$.
- Notice that $\arg(z)$ and $\mathrm{Arg}(z)$ are not defined if $z = (0, 0)$.



Example

Let $z = (\sqrt{3}, 1)$.
Then $\mathrm{Re}(z) = \sqrt{3}$, $\mathrm{Im}(z) = 1$, and
$|z| = \sqrt{(\sqrt{3})^2 + 1^2} = \sqrt{4} = 2$.
And $\mathrm{Arg}(z) = \arctan\left(\frac{\mathrm{Im}(z)}{\mathrm{Re}(z)}\right) =$
$\arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}$, and
$\arg(z) = \frac{\pi}{6} + 2\pi k$, $k \in \mathbb{Z}$.



NTNU
Norwegian University of
Science and Technology

How to compute the argument of a complex number

The argument of a complex number z can be computed using trigonometric identities, $\mathrm{Re}(z)$, $\mathrm{Im}(z)$, $|z|$ and arccos, arcsin and arctan.

- If $\mathrm{Re}(z) > 0$, then
  $\mathrm{Arg}(z) = \arctan\left(\frac{\mathrm{Im}(z)}{\mathrm{Re}(z)}\right) = \arcsin\left(\frac{\mathrm{Im}(z)}{|z|}\right)$.
- If $\mathrm{Im}(z) \geq 0$, then $\mathrm{Arg}(z) = \arccos\left(\frac{\mathrm{Re}(z)}{|z|}\right)$.
- $\mathrm{Arg}(z) = \mathrm{Arg}(-z) \pm \pi$.

NTNU
Norwegian University of
Science and Technology

Example

Let $z = \left(\frac{1}{4}, \frac{-\sqrt{3}}{4}\right)$.

Then $\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{-\sqrt{3}/4}{1/4}\right) =$

$\arctan(-\sqrt{3}) = -\arctan(\sqrt{3}) = \frac{-\pi}{3}$.

We could also compute $\text{Arg}(z)$ in the following way:

$$\text{Arg}(z) = \arcsin\left(\frac{\text{Im}(z)}{|z|}\right) = \arcsin\left(\frac{-\sqrt{3}/4}{\sqrt{(1/4)^2+(-\sqrt{3}/4)^2}}\right) =$$

$$\arcsin\left(\frac{-\sqrt{3}}{2}\right) = -\arcsin\left(\frac{\sqrt{3}}{2}\right) = \frac{-\pi}{3}.$$

NTNU
Norwegian University of
Science and Technology

Example

Let $z = (-1, 1)$.

Then $\text{Arg}(z) = \arccos\left(\frac{\text{Re}(z)}{|z|}\right) = \arccos\left(\frac{-1}{\sqrt{(-1)^2+1^2}}\right) =$

$\arccos(\frac{-1}{\sqrt{2}}) = \frac{3\pi}{4}$.

We could also compute $\text{Arg}(z)$ in the following way:

$\text{Arg}(z) = \text{Arg}(-z) + \pi = \arctan\left(\frac{\text{Im}(-z)}{\text{Re}(-z)}\right) + \pi =$

$\arctan\left(\frac{-1}{1}\right) + \pi = -\arctan(1) + \pi = -\frac{\pi}{4} + \pi = \frac{3\pi}{4}$.

Example

Let $z = (-2, -2)$.
Then $\text{Arg}(z) = \text{Arg}(-z) - \pi = \arctan\left(\frac{\text{Im}(-z)}{\text{Re}(-z)}\right) + \pi =$
$\arctan\left(\frac{-2}{-2}\right) - \pi = \arctan(1) - \pi = \frac{\pi}{4} - \pi = \frac{-3\pi}{4}$.

NTNU
Norwegian University of
Science and Technology

How do we add real numbers?

If a and b are real numbers, then we get $a + b$ by taking the line from 0 to b and move it such that it starts at a.



NTNU
Norwegian University of
Science and Technology

How do we add complex numbers?

If z and w are complex numbers, then we get $z + w$ by taking the line from 0 to w and move it such that it starts at z.



NTNU
Norwegian University of
Science and Technology

Example

What is $(2,1)+(1,2)$?



NTNU
Norwegian University of
Science and Technology

Addition of complex numbers

If z_1, z_2 and z_3 are complex numbers, then

- $\mathrm{Re}(z_1 + z_2) = \mathrm{Re}(z_1) + \mathrm{Re}(z_2)$ and $\mathrm{Im}(z_1 + z_2) = \mathrm{Im}(z_1) + \mathrm{Im}(z_2)$,
- $z_1 + z_2 = z_2 + z_1$,
- $(z_1 + z_2) + z_3 = z_1 + (z_2 + z_3)$,
- $|z_1 + z_2| \leq |z_1| + |z_2|$.

NTNU
Norwegian University of
Science and Technology

How do we multiply real numbers?

If a and b are real numbers and $b > 0$, then we get ab by scaling the line from 0 to a by a factor of b.



How do we multiply real numbers?

If a and b are real numbers and $b < 0$, then we get ab by rotating the line from 0 to a by 180° degrees and then scale it by a factor of $|b|$.



NTNU
Norwegian University of
Science and Technology

How do we multiply complex numbers?

If z and w are real numbers, then we get zw by rotating the line from 0 to z by $\arg(w)$ and then scale it by a factor of $|w|$.



NTNU
Norwegian University of Science and Technology

Multiplication of complex numbers

If z_1, z_2 and z_3 are complex numbers, then

- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$ and $|z_1 z_2| = |z_1||z_2|$,
- $z_1 z_2 = z_2 z_1$,
- $(z_1 z_2) z_3 = z_1 (z_2 z_3)$,
- $z_1 (z_2 + z_3) = z_1 z_2 + z_1 z_3$.

NTNU
Norwegian University of
Science and Technology

Example

Let $z_1 = (\sqrt{3}, 1)$ and $z_2 = (\frac{1}{4}, \frac{-\sqrt{3}}{4})$.
We have shown that $|z_1| = 2$, $\text{Arg}(z_1) = \frac{\pi}{6}$, $|z_2| = 1/2$ and $\text{Arg}(z_2) = \frac{-\pi}{3}$.
It follows that $|z_1 z_2| = |z_1||z_2| = 2(1/2) = 1$ and $\arg(z_1 z_2) = \text{Arg}(z_1) + \text{Arg}(z_2) = \frac{\pi}{6} - \frac{\pi}{3} = \frac{-\pi}{6}$.
So $z_1 z_2 = (|z_1 z_2| \cos(\arg(z_1 z_2)), |z_1 z_2| \sin(z_1 z_2)) = (\cos(-\pi/6), \sin(-\pi/6)) = \left(\frac{\sqrt{3}}{2}, \frac{-1}{2}\right)$.

NTNU
Norwegian University of
Science and Technology

Powers of complex numbers

If z is a complex number and n is a positive integer, then $\arg(z^n) = n\arg(z)$ and $|z^n| = |z|^n$.

NTNU
Norwegian University of
Science and Technology

Example

Let $i = (0, 1)$.
Then $\text{Arg}(i) = \frac{\pi}{2}$ and $|i| = 1$.
It follows that $\text{Arg}(i^2) = \pi$ and $|i^2| = 1$.
So $i^2 = (\cos(\pi), \sin(\pi)) = (1, 0)$.

Notation

- When a is a real number, then we identify $(a, 0)$ with a.
- Let $i = (0, 1)$.
- Then every complex number (a, b) can be written as $a + bi$.
- $\mathrm{Re}(a + bi) = a$, $\mathrm{Im}(a + bi) = b$ and $|a + bi| = \sqrt{a^2 + b^2}$.
- $(a + bi) + (c + di) = (a + c) + (b + d)i$.
- $i^2 = -1$
- $(a + bi)(c + di) = ac - bd + (ad + bc)i$.

NTNU
Norwegian University of
Science and Technology

Complex conjugation

- When $z = a + bi$ is a complex number, then the number $a - bi$ is called the *conjugate* of z and is denoted by $\overline{z}$.
- $\mathrm{Re}(\overline{z}) = \mathrm{Re}(z)$ and $\mathrm{Im}(\overline{z}) = -\mathrm{Im}(z)$.
- We get $\overline{z}$ by reflecting z in the real line.
- $|\overline{z}| = |z|$ and $\arg(\overline{z}) = -\arg(z)$.



NTNU
Norwegian University of
Science and Technology

Complex conjugation

- $z = \overline{z}$ if and only if $\mathrm{Im}(z) = 0$.
- $z = -\overline{z}$ if and only if $\mathrm{Re}(z) = 0$.
- $\overline{z + w} = \overline{z} + \overline{w}$.
- $\overline{zw} = \overline{z}\,\overline{w}$.
- $z\overline{z} = |z|^2$.
- $\dfrac{z}{w} = \dfrac{z\overline{w}}{w\overline{w}} = \dfrac{z\overline{w}}{|w|^2}$.

NTNU
Norwegian University of
Science and Technology

Example

$$\frac{2+1}{3-1} = \frac{(2+i)(3+i)}{(3-i)(3+i)} = \frac{6-1+2i+3i}{9+1} = \frac{5+5i}{10} = \frac{1}{2} + \frac{1}{2}i.$$