



NTNU  
Norwegian University of  
Science and Technology

**TMA4115 - Calculus 3**  
**Lecture 2, Jan 17**

Toke Meier Carlsen  
Norwegian University of Science and Technology  
Spring 2013

# Course web page

Information about the course can be found at  
<http://wiki.math.ntnu.no/tma4115/2013v>.



NTNU  
Norwegian University of  
Science and Technology

# Review of yesterday's lecture



NTNU  
Norwegian University of  
Science and Technology

# Review of yesterday's lecture

Yesterday we introduced *complex numbers* and studied



NTNU  
Norwegian University of  
Science and Technology

# Review of yesterday's lecture

Yesterday we introduced *complex numbers* and studied

- the *real part*, the *imaginary part*, the *absolute value* (or *modulus*), and the *argument* of a complex number,



# Review of yesterday's lecture

Yesterday we introduced *complex numbers* and studied

- the *real part*, the *imaginary part*, the *absolute value* (or *modulus*), and the *argument* of a complex number,
- addition and multiplication of complex numbers,



# Review of yesterday's lecture

Yesterday we introduced *complex numbers* and studied

- the *real part*, the *imaginary part*, the *absolute value* (or *modulus*), and the *argument* of a complex number,
- addition and multiplication of complex numbers,
- and *complex conjugation*.



# Today's lecture



NTNU  
Norwegian University of  
Science and Technology



# Today's lecture

Today we will study



NTNU  
Norwegian University of  
Science and Technology

# Today's lecture

Today we will study

- *polar representation* of complex numbers,



NTNU  
Norwegian University of  
Science and Technology

# Today's lecture

Today we will study

- *polar representation* of complex numbers,
- *de Moivre's Theorem*,



NTNU  
Norwegian University of  
Science and Technology

# Today's lecture

Today we will study

- *polar representation* of complex numbers,
- *de Moivre's Theorem*,
- how complex numbers can be used to derive trigonometric identities,



NTNU  
Norwegian University of  
Science and Technology

# Today's lecture

Today we will study

- *polar representation* of complex numbers,
- *de Moivre's Theorem*,
- how complex numbers can be used to derive trigonometric identities,
- roots of complex numbers.




# Complex numbers

- A complex number is a number which can be written as  $a + ib$  where  $a$  and  $b$  are real numbers and  $i$  satisfies  $i^2 = -1$ .




# Complex numbers

- A complex number is a number which can be written as  $a + ib$  where  $a$  and  $b$  are real numbers and  $i$  satisfies  $i^2 = -1$ .
- A complex number  $z = a + ib$  can be represented as the point  $(a, b)$  in the plane.



# Complex numbers


- A complex number is a number which can be written as  $a + ib$  where  $a$  and  $b$  are real numbers and  $i$  satisfies  $i^2 = -1$ .
- A complex number  $z = a + ib$  can be represented as the point  $(a, b)$  in the plane.
- If  $z = a + ib$ , then  $a$  is called the *real part* of  $z$  and is denoted by  $\text{Re}(z)$ .






# Complex numbers

- A complex number is a number which can be written as  $a + ib$  where  $a$  and  $b$  are real numbers and  $i$  satisfies  $i^2 = -1$ .
- A complex number  $z = a + ib$  can be represented as the point  $(a, b)$  in the plane.
- If  $z = a + ib$ , then  $a$  is called the *real part* of  $z$  and is denoted by  $\operatorname{Re}(z)$ .
- $b$  is called the *imaginary part* of  $z$  and is denoted by  $\operatorname{Im}(z)$ .




# Complex numbers

- If  $z = a + ib$ , then the length  $\sqrt{a^2 + b^2}$  of the line from  $(0, 0)$  to  $(a, b)$  is called the *modulus* or the *absolute value* of  $z$  and is denoted by  $|z|$ .



# Complex numbers

- If  $z = a + ib$ , then the length  $\sqrt{a^2 + b^2}$  of the line from  $(0, 0)$  to  $(a, b)$  is called the *modulus* or the *absolute value* of  $z$  and is denoted by  $|z|$ .
- The angle between the line through  $(0, 0)$  and  $(a, b)$  and the positive part of the real axis is called the *argument* of  $z$  and is denoted by  $\arg(z)$ .



# Complex numbers

- $\arg(z)$  is not unique. If  $\theta = \arg(z)$ , then also  $\theta + 2\pi = \arg(z)$ . If we want to be precise, then  $\arg(z)$  is really the set of all angles  $\theta$  which satisfies that if we rotate the positive part of the real axis by  $\theta$ , then it lands on the line through  $(0, 0)$  and  $(a, b)$ .



# Complex numbers

- $\arg(z)$  is not unique. If  $\theta = \arg(z)$ , then also  $\theta + 2\pi = \arg(z)$ . If we want to be precise, then  $\arg(z)$  is really the set of all angles  $\theta$  which satisfies that if we rotate the positive part of the real axis by  $\theta$ , then it lands on the line through  $(0, 0)$  and  $(a, b)$ .
- The unique value of  $\arg(z)$  in the interval  $(-\pi, \pi]$  is called the *principal argument* of  $z$  and is denoted by  $\text{Arg}(z)$ .



# Complex numbers

- $\arg(z)$  is not unique. If  $\theta = \arg(z)$ , then also  $\theta + 2\pi = \arg(z)$ . If we want to be precise, then  $\arg(z)$  is really the set of all angles  $\theta$  which satisfies that if we rotate the positive part of the real axis by  $\theta$ , then it lands on the line through  $(0, 0)$  and  $(a, b)$ .
- The unique value of  $\arg(z)$  in the interval  $(-\pi, \pi]$  is called the *principal argument* of  $z$  and is denoted by  $\text{Arg}(z)$ .
- Notice that  $\arg(z)$  and  $\text{Arg}(z)$  are not defined if  $z = (0, 0)$ .



# Addition of complex numbers



NTNU  
Norwegian University of  
Science and Technology

# Addition of complex numbers

- If  $z_1 = a + bi$  and  $z_2 = c + di$ , then  $z_1 + z_2 = (a + c) + (b + d)i$ .
- $\operatorname{Re}(z_1 + z_2) = \operatorname{Re}(z_1) + \operatorname{Re}(z_2)$  and  $\operatorname{Im}(z_1 + z_2) = \operatorname{Im}(z_1) + \operatorname{Im}(z_2)$ .





# Addition of complex numbers

- If  $z_1 = a + bi$  and  $z_2 = c + di$ , then  $z_1 + z_2 = (a + c) + (b + d)i$ .
- $\operatorname{Re}(z_1 + z_2) = \operatorname{Re}(z_1) + \operatorname{Re}(z_2)$  and  $\operatorname{Im}(z_1 + z_2) = \operatorname{Im}(z_1) + \operatorname{Im}(z_2)$ .
- $z_1 + z_2 = z_2 + z_1$ .
- $(z_1 + z_2) + z_3 = z_1 + (z_2 + z_3)$ .
- $|z_1 + z_2| \leq |z_1| + |z_2|$ .



# Multiplication of complex numbers



NTNU  
Norwegian University of  
Science and Technology

# Multiplication of complex numbers

- If  $z_1 = a + bi$  and  $z_2 = c + di$ , then
$$z_1 z_2 = (a + bi)(c + di) = ac + adi + bci + bdi^2 = (ac - bd) + (ad + bc)i.$$



# Multiplication of complex numbers

- If  $z_1 = a + bi$  and  $z_2 = c + di$ , then
$$z_1 z_2 = (a + bi)(c + di) = ac + adi + bci + bdi^2 = (ac - bd) + (ad + bc)i.$$
- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$  and  $|z_1 z_2| = |z_1| |z_2|$ .



# Multiplication of complex numbers

- If  $z_1 = a + bi$  and  $z_2 = c + di$ , then
$$z_1 z_2 = (a + bi)(c + di) = ac + adi + bci + bdi^2 = (ac - bd) + (ad + bc)i.$$
- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$  and  $|z_1 z_2| = |z_1| |z_2|$ .
- $z_1 z_2 = z_2 z_1$ .
- $(z_1 z_2) z_3 = z_1 (z_2 z_3)$ .
- $z_1 (z_2 + z_3) = z_1 z_2 + z_1 z_3$ .



# Complex conjugation



NTNU  
Norwegian University of  
Science and Technology

# Complex conjugation

- When  $z = a + bi$  is a complex number, then the number  $a - bi$  is called the *conjugate* of  $z$  and is denoted by  $\bar{z}$ .



# Complex conjugation


- When  $z = a + bi$  is a complex number, then the number  $a - bi$  is called the *conjugate* of  $z$  and is denoted by  $\bar{z}$ .
- $\operatorname{Re}(\bar{z}) = \operatorname{Re}(z)$  and  $\operatorname{Im}(\bar{z}) = -\operatorname{Im}(z)$ .






# Complex conjugation

- When  $z = a + bi$  is a complex number, then the number  $a - bi$  is called the *conjugate* of  $z$  and is denoted by  $\bar{z}$ .
- $\operatorname{Re}(\bar{z}) = \operatorname{Re}(z)$  and  $\operatorname{Im}(\bar{z}) = -\operatorname{Im}(z)$ .
- We get  $\bar{z}$  by reflecting  $z$  in the real line.




# Complex conjugation

- When  $z = a + bi$  is a complex number, then the number  $a - bi$  is called the *conjugate* of  $z$  and is denoted by  $\bar{z}$ .
- $\operatorname{Re}(\bar{z}) = \operatorname{Re}(z)$  and  $\operatorname{Im}(\bar{z}) = -\operatorname{Im}(z)$ .
- We get  $\bar{z}$  by reflecting  $z$  in the real line.



# Complex conjugation

- When  $z = a + bi$  is a complex number, then the number  $a - bi$  is called the *conjugate* of  $z$  and is denoted by  $\bar{z}$ .
- $\operatorname{Re}(\bar{z}) = \operatorname{Re}(z)$  and  $\operatorname{Im}(\bar{z}) = -\operatorname{Im}(z)$ .
- We get  $\bar{z}$  by reflecting  $z$  in the real line.
- $|\bar{z}| = |z|$  and  $\arg(\bar{z}) = -\arg(z)$ .



# Complex conjugation

- $z = \bar{z}$  if and only if  $\text{Im}(z) = 0$ .
- $z = -\bar{z}$  if and only if  $\text{Re}(z) = 0$ .
- $\overline{z + w} = \bar{z} + \bar{w}$ .
- $\overline{zw} = \bar{z} \bar{w}$ .
- $z\bar{z} = (a+bi)(a-bi) = a^2 - abi + abi - b^2i^2 = a^2 + b^2 = |z|^2$ .



# Division of complex numbers



NTNU  
Norwegian University of  
Science and Technology

# Division of complex numbers

- If  $z$  and  $w$  are complex numbers and  $w \neq 0$ , then

$$\frac{z}{w} = \frac{z\bar{w}}{w\bar{w}} = \frac{z\bar{w}}{|w|^2}.$$



# Division of complex numbers

- If  $z$  and  $w$  are complex numbers and  $w \neq 0$ , then

$$\frac{z}{w} = \frac{z\bar{w}}{w\bar{w}} = \frac{z\bar{w}}{|w|^2}.$$

- $\frac{a + bi}{c + di} = \frac{(a + bi)(c - di)}{c^2 + d^2} = \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i.$



# Division of complex numbers

- If  $z$  and  $w$  are complex numbers and  $w \neq 0$ , then

$$\frac{z}{w} = \frac{z\bar{w}}{w\bar{w}} = \frac{z\bar{w}}{|w|^2}.$$

- $$\frac{a + bi}{c + di} = \frac{(a + bi)(c - di)}{c^2 + d^2} = \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i.$$

- $$\arg\left(\frac{z}{w}\right) = \arg(z) - \arg(w) \text{ and } \left|\frac{z}{w}\right| = \frac{|z|}{|w|}.$$





# Example



NTNU  
Norwegian University of  
Science and Technology

# Example

$$\frac{1+i}{2-i} = \frac{(1+i)(2+i)}{(2-i)(2+i)} = \frac{2-1+3i}{5} = \frac{1}{5} + \frac{3}{5}i.$$




# Polar representation



NTNU  
Norwegian University of  
Science and Technology


# Polar representation

- Every complex number  $z$  can be written on the form  $r(\cos \theta + i \sin \theta)$  where  $r$  and  $\theta$  are real numbers and  $r \geq 0$ . This is called the *polar form* of  $z$ .



# Polar representation

- Every complex number  $z$  can be written on the form  $r(\cos \theta + i \sin \theta)$  where  $r$  and  $\theta$  are real numbers and  $r \geq 0$ . This is called the *polar form* of  $z$ .
- Notice that if  $z = r(\cos \theta + i \sin \theta)$ , then  $|z| = r$  and  $\arg(z) = \theta$ .



# Multiplication and division using polar representations



NTNU  
Norwegian University of  
Science and Technology

# Multiplication and division using polar representations

Recall that

- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$  and  $|z_1 z_2| = |z_1| |z_2|$ , and that
- $\arg\left(\frac{z}{w}\right) = \arg(z) - \arg(w)$  and  $\left|\frac{z}{w}\right| = \frac{|z|}{|w|}$ .



# Multiplication and division using polar representations

Recall that

- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$  and  $|z_1 z_2| = |z_1| |z_2|$ , and that
- $\arg\left(\frac{z}{w}\right) = \arg(z) - \arg(w)$  and  $\left|\frac{z}{w}\right| = \frac{|z|}{|w|}$ .

It follows that if  $z = r_1(\cos \theta_1 + i \sin \theta_1)$  and  $z_2 = r_2(\cos \theta_2 + i \sin \theta_2)$ , then





# Multiplication and division using polar representations

Recall that

- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$  and  $|z_1 z_2| = |z_1| |z_2|$ , and that
- $\arg\left(\frac{z}{w}\right) = \arg(z) - \arg(w)$  and  $\left|\frac{z}{w}\right| = \frac{|z|}{|w|}$ .

It follows that if  $z = r_1(\cos \theta_1 + i \sin \theta_1)$  and  $z_2 = r_2(\cos \theta_2 + i \sin \theta_2)$ , then

- $z_1 z_2 = r_1 r_2 (\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2))$  and



# Multiplication and division using polar representations

Recall that

- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$  and  $|z_1 z_2| = |z_1| |z_2|$ , and that
- $\arg\left(\frac{z}{w}\right) = \arg(z) - \arg(w)$  and  $\left|\frac{z}{w}\right| = \frac{|z|}{|w|}$ .

It follows that if  $z = r_1(\cos \theta_1 + i \sin \theta_1)$  and  $z_2 = r_2(\cos \theta_2 + i \sin \theta_2)$ , then

- $z_1 z_2 = r_1 r_2 (\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2))$  and
- $\frac{z_1}{z_2} = \frac{r_1}{r_2} (\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2))$  (provided  $r_2 \neq 0$ ).



# Exercise 46, page xviii

Express each of the complex numbers  $z = 3 + i\sqrt{3}$  and  $w = -1 + i\sqrt{3}$  in polar form. Use these expressions to calculate  $zw$  and  $z/w$ .



# Solution



NTNU  
Norwegian University of  
Science and Technology

# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2,$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2, \text{ and}$$

$$\text{Arg}(w) = \arccos\left(\frac{\text{Re}(w)}{|w|}\right) = \arccos\left(\frac{-1}{2}\right) = \frac{2\pi}{3},$$





# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2, \text{ and}$$

$$\text{Arg}(w) = \arccos\left(\frac{\text{Re}(w)}{|w|}\right) = \arccos\left(\frac{-1}{2}\right) = \frac{2\pi}{3},$$

$$\text{so } |zw| = |z||w| = (\sqrt{12})2 = 4\sqrt{3},$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2, \text{ and}$$

$$\text{Arg}(w) = \arccos\left(\frac{\text{Re}(w)}{|w|}\right) = \arccos\left(\frac{-1}{2}\right) = \frac{2\pi}{3},$$

$$\text{so } |zw| = |z||w| = (\sqrt{12})2 = 4\sqrt{3},$$

$$\text{arg}(zw) = \text{Arg}(z) + \text{Arg}(w) = \frac{\pi}{6} + \frac{2\pi}{3} = \frac{5\pi}{6},$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2, \text{ and}$$

$$\text{Arg}(w) = \arccos\left(\frac{\text{Re}(w)}{|w|}\right) = \arccos\left(\frac{-1}{2}\right) = \frac{2\pi}{3},$$

$$\text{so } |zw| = |z||w| = (\sqrt{12})2 = 4\sqrt{3},$$

$$\arg(zw) = \text{Arg}(z) + \text{Arg}(w) = \frac{\pi}{6} + \frac{2\pi}{3} = \frac{5\pi}{6},$$

$$|z/w| = |z|/|w| = \sqrt{12}/2 = \sqrt{3},$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2, \text{ and}$$

$$\text{Arg}(w) = \arccos\left(\frac{\text{Re}(w)}{|w|}\right) = \arccos\left(\frac{-1}{2}\right) = \frac{2\pi}{3},$$

$$\text{so } |zw| = |z||w| = (\sqrt{12})2 = 4\sqrt{3},$$

$$\arg(zw) = \text{Arg}(z) + \text{Arg}(w) = \frac{\pi}{6} + \frac{2\pi}{3} = \frac{5\pi}{6},$$

$$|z/w| = |z|/|w| = \sqrt{12}/2 = \sqrt{3}, \text{ and}$$

$$\arg(z/w) = \text{Arg}(z) - \text{Arg}(w) = \frac{\pi}{6} - \frac{2\pi}{3} = \frac{-\pi}{2},$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2, \text{ and}$$

$$\text{Arg}(w) = \arccos\left(\frac{\text{Re}(w)}{|w|}\right) = \arccos\left(\frac{-1}{2}\right) = \frac{2\pi}{3},$$

$$\text{so } |zw| = |z||w| = (\sqrt{12})2 = 4\sqrt{3},$$

$$\arg(zw) = \text{Arg}(z) + \text{Arg}(w) = \frac{\pi}{6} + \frac{2\pi}{3} = \frac{5\pi}{6},$$

$$|z/w| = |z|/|w| = \sqrt{12}/2 = \sqrt{3}, \text{ and}$$

$$\arg(z/w) = \text{Arg}(z) - \text{Arg}(w) = \frac{\pi}{6} - \frac{2\pi}{3} = \frac{-\pi}{2}, \text{ from which it}$$

$$\text{follows that } zw = 4\sqrt{3}(\cos(5\pi/6) + i \sin(5\pi/6)) =$$

$$4\sqrt{3}\left(\frac{-\sqrt{3}}{2} + \frac{1}{2}i\right) = -6 + 2\sqrt{3}i,$$



# Solution

$$|z| = \sqrt{3^2 + (\sqrt{3})^2} = \sqrt{12},$$

$$\text{Arg}(z) = \arctan\left(\frac{\text{Im}(z)}{\text{Re}(z)}\right) = \arctan\left(\frac{\sqrt{3}}{3}\right) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6},$$

$$|w| = \sqrt{(-1)^2 + (\sqrt{3})^2} = \sqrt{4} = 2, \text{ and}$$

$$\text{Arg}(w) = \arccos\left(\frac{\text{Re}(w)}{|w|}\right) = \arccos\left(\frac{-1}{2}\right) = \frac{2\pi}{3},$$

$$\text{so } |zw| = |z||w| = (\sqrt{12})2 = 4\sqrt{3},$$

$$\arg(zw) = \text{Arg}(z) + \text{Arg}(w) = \frac{\pi}{6} + \frac{2\pi}{3} = \frac{5\pi}{6},$$

$$|z/w| = |z|/|w| = \sqrt{12}/2 = \sqrt{3}, \text{ and}$$

$$\arg(z/w) = \text{Arg}(z) - \text{Arg}(w) = \frac{\pi}{6} - \frac{2\pi}{3} = \frac{-\pi}{2}, \text{ from which it}$$

$$\text{follows that } zw = 4\sqrt{3}(\cos(5\pi/6) + i \sin(5\pi/6)) =$$

$$4\sqrt{3}\left(\frac{-\sqrt{3}}{2} + \frac{1}{2}i\right) = -6 + 2\sqrt{3}i, \text{ and } z/w =$$

$$\sqrt{3}(\cos(-\pi/2) + i \sin(-\pi/2)) = -\sqrt{3}i.$$



# Complex numbers and trigonometric identities



NTNU  
Norwegian University of  
Science and Technology

# Complex numbers and trigonometric identities

Polar representations of complex numbers can be used to derive trigonometric identities like

- $\cos(\theta_1 + \theta_2) = \cos(\theta_1) \cos(\theta_2) - \sin(\theta_1) \sin(\theta_2)$
- $\sin(\theta_1 + \theta_2) = \cos(\theta_1) \sin(\theta_2) + \sin(\theta_1) \cos(\theta_2)$





# Proof



NTNU  
Norwegian University of  
Science and Technology

# Proof

We have that

$$\begin{aligned}\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) &= (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2) \\ &= \cos \theta_1 \cos \theta_2 + i \cos \theta_1 \sin \theta_2 + i \sin \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2 \\ &= \cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2 + i(\cos \theta_1 \sin \theta_2 + \sin \theta_1 \cos \theta_2)\end{aligned}$$



# Proof

We have that

$$\begin{aligned}\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) &= (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2) \\ &= \cos \theta_1 \cos \theta_2 + i \cos \theta_1 \sin \theta_2 + i \sin \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2 \\ &= \cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2 + i(\cos \theta_1 \sin \theta_2 + \sin \theta_1 \cos \theta_2)\end{aligned}$$

from which it follows that

$$\begin{aligned}\cos(\theta_1 + \theta_2) &= \cos(\theta_1) \cos(\theta_2) - \sin(\theta_1) \sin(\theta_2) \text{ and} \\ \sin(\theta_1 + \theta_2) &= \cos(\theta_1) \sin(\theta_2) + \sin(\theta_1) \cos(\theta_2).\end{aligned}$$



# de Moivre's Theorem

- Recall that  $\arg(zw) = \arg(z) + \arg(w)$  and  $|zw| = |z||w|$ .



# de Moivre's Theorem

- Recall that  $\arg(zw) = \arg(z) + \arg(w)$  and  $|zw| = |z||w|$ .
- It follows that
$$(r(\cos \theta + i \sin \theta))^n = r^n(\cos(n\theta) + i \sin(n\theta)).$$



# de Moivre's Theorem

- Recall that  $\arg(zw) = \arg(z) + \arg(w)$  and  $|zw| = |z||w|$ .
- It follows that
$$(r(\cos \theta + i \sin \theta))^n = r^n(\cos(n\theta) + i \sin(n\theta)).$$
- We have in particular that

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta).$$



# de Moivre's Theorem

- Recall that  $\arg(zw) = \arg(z) + \arg(w)$  and  $|zw| = |z||w|$ .
- It follows that
$$(r(\cos \theta + i \sin \theta))^n = r^n(\cos(n\theta) + i \sin(n\theta)).$$
- We have in particular that

$$(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta).$$

- This formula is known as *de Moivre's formula*.



# Complex numbers and trigonometric identities



NTNU  
Norwegian University of  
Science and Technology



# Complex numbers and trigonometric identities

Polar representations of complex numbers can be used to derive trigonometric identities like

- $\cos(\theta_1 + \theta_2) = \cos(\theta_1) \cos(\theta_2) - \sin(\theta_1) \sin(\theta_2)$
- $\sin(\theta_1 + \theta_2) = \cos(\theta_1) \sin(\theta_2) + \sin(\theta_1) \cos(\theta_2)$



# Complex numbers and trigonometric identities

Polar representations of complex numbers can be used to derive trigonometric identities like

- $\cos(\theta_1 + \theta_2) = \cos(\theta_1) \cos(\theta_2) - \sin(\theta_1) \sin(\theta_2)$
- $\sin(\theta_1 + \theta_2) = \cos(\theta_1) \sin(\theta_2) + \sin(\theta_1) \cos(\theta_2)$
- $\cos(3\theta) = \cos^3(\theta) - 3 \cos(\theta) \sin^2(\theta)$
- $\sin(3\theta) = 3 \cos^2(\theta) \sin(\theta) - \sin^3(\theta)$



# Proof



NTNU  
Norwegian University of  
Science and Technology

# Proof

It follows from de Moivre's formula and the binomial formula that

$$\begin{aligned}\cos(3\theta) + i \sin(3\theta) &= \cos^3(\theta) + 3i \cos^2(\theta) \sin(\theta) + 3i^2 \cos(\theta) \sin^2(\theta) + i^3 \sin^3(\theta) \\ &= \cos^3(\theta) - 3 \cos(\theta) \sin^2(\theta) + i(3 \cos^2(\theta) \sin(\theta) - \sin^3(\theta))\end{aligned}$$



# Proof

It follows from de Moivre's formula and the binomial formula that

$$\begin{aligned}\cos(3\theta) + i \sin(3\theta) &= \cos^3(\theta) + 3i \cos^2(\theta) \sin(\theta) + 3i^2 \cos(\theta) \sin^2(\theta) + i^3 \sin^3(\theta) \\ &= \cos^3(\theta) - 3 \cos(\theta) \sin^2(\theta) + i(3 \cos^2(\theta) \sin(\theta) - \sin^3(\theta))\end{aligned}$$

from which it follows that  $\cos(3\theta) = \cos^3(\theta) - 3 \cos(\theta) \sin^2(\theta)$  and  $\sin(3\theta) = 3 \cos^2(\theta) \sin(\theta) - \sin^3(\theta)$ .



# Roots of complex numbers



NTNU  
Norwegian University of  
Science and Technology

# Roots of complex numbers

- If  $z$  is a complex number and  $n$  is a positive integer, then an  $n$ 'th root of  $z$  is a complex number  $w$  satisfying  $w^n = z$ .



# Roots of complex numbers

- If  $z$  is a complex number and  $n$  is a positive integer, then an  $n$ 'th root of  $z$  is a complex number  $w$  satisfying  $w^n = z$ .
- Every complex number different from 0 has  $n$  different  $n$ 'th roots.





# Example

Let us find the 3 cube roots (3rd roots) of  $z = -2 - 2i$ .



# Example

Let us find the 3 cube roots (3rd roots) of  $z = -2 - 2i$ .

$$|z| = \sqrt{(-2)^2 + (-2)^2} = \sqrt{8} \text{ and } \text{Arg}(z) = \frac{-3\pi}{4},$$



# Example

Let us find the 3 cube roots (3rd roots) of  $z = -2 - 2i$ .

$|z| = \sqrt{(-2)^2 + (-2)^2} = \sqrt{8}$  and  $\text{Arg}(z) = \frac{-3\pi}{4}$ , so it follows from de Moivre's formula that

$$w^3 = z \iff |w|^3 = |z| = \sqrt{8} \text{ and}$$

$$3 \arg(w) = \arg(z) = \frac{-3\pi}{4} + 2\pi k, \quad k \in \mathbb{Z}$$

$$\iff |w| = \sqrt[3]{\sqrt{8}} = 8^{1/6} = \sqrt{2} \text{ and}$$

$$\arg w = \frac{-\pi}{4} + \frac{2\pi k}{3}, \quad k \in \mathbb{Z}.$$



# Example

So the 3 cube roots of  $z = -2 - 2i$  are

$$\begin{aligned}w_0 &= \sqrt{2}(\cos(-\pi/4) + i \sin(-\pi/4)) \\ &= \sqrt{2} \left( \frac{1}{\sqrt{2}} - \frac{1}{\sqrt{2}}i \right) = 1 - i\end{aligned}$$

$$\begin{aligned}w_1 &= \sqrt{2}(\cos(5\pi/12) + i \sin(5\pi/12)) \\ &= \sqrt{2} \left( \frac{\sqrt{3}-1}{2\sqrt{2}} + i \frac{1+\sqrt{3}}{2\sqrt{2}}i \right) = \frac{\sqrt{3}-1}{2} + i \frac{1+\sqrt{3}}{2}\end{aligned}$$

$$\begin{aligned}w_2 &= \sqrt{2}(\cos(13\pi/12) + i \sin(13\pi/12)) \\ &= \sqrt{2} \left( -\frac{1+\sqrt{3}}{2\sqrt{2}} + i \frac{\sqrt{3}-1}{2\sqrt{2}}i \right) = -\frac{1+\sqrt{3}}{2} + i \frac{\sqrt{3}-1}{2}\end{aligned}$$



# Roots of complex numbers

If  $z$  is a complex number different from 0 and  $n$  is a positive integer, then the  $n$ 'th roots of  $z$  are

$$w_k = |z|^{1/n} \left( \cos \left( \frac{\arg(z) + 2\pi k}{n} \right) + i \sin \left( \frac{\arg(z) + 2\pi k}{n} \right) \right)$$

where  $k = 0, 1, \dots, n - 1$ .



# Roots of unity

- If  $n$  is a positive integer, then a  $n$ 'th root of 1 is called a  *$n$ 'th root of unity*.



# Roots of unity

- If  $n$  is a positive integer, then a  $n$ 'th root of 1 is called a  *$n$ 'th root of unity*.
- The  $n$   $n$ 'th roots of unity are

$$u_n = \cos\left(\frac{2\pi k}{n}\right) + i \sin\left(\frac{2\pi k}{n}\right)$$

for  $k = 0, 1, \dots, n - 1$ .




# The 6<sup>th</sup> root of unity



NTNU  
Norwegian University of  
Science and Technology



# The 6<sup>th</sup> root of unity



# Roots of complex numbers

If  $z$  is a complex number different from 0,  $n$  is a positive integer and  $w$  is an  $n$ 'th root of  $z$ , then the other  $n - 1$   $n$ 'th roots of  $z$  are  $wu_1, wu_2, \dots, wu_{n-1}$  where  $u_1, u_2, \dots, u_{n-1}$  are the  $n - 1$   $n$ 'th roots of unity different from 1.



# Problem 1 from the exam from August 2012

Write all of the solutions of  $z^3 = 1$  in the form  $z = x + iy$ .  
Write the solutions of  $z^3 = \frac{-3+i}{\sqrt{2}(2+i)}$  in the form  $z = x + iy$  and draw the solutions in the complex plane.



# Solution



NTNU  
Norwegian University of  
Science and Technology

# Solution

The solutions of  $z^3 = 1$  are the 3 cube roots of unity

$$u_0 = \cos(0) + i \sin(0) = 1$$

$$u_1 = \cos(2\pi/3) + i \sin(2\pi/3) = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$$

$$u_2 = \cos(4\pi/3) + i \sin(4\pi/3) = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$$



# Solution

The solutions of  $z^3 = 1$  are the 3 cube roots of unity

$$u_0 = \cos(0) + i \sin(0) = 1$$

$$u_1 = \cos(2\pi/3) + i \sin(2\pi/3) = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$$

$$u_2 = \cos(4\pi/3) + i \sin(4\pi/3) = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$$

$$\begin{aligned} \frac{-3+i}{\sqrt{2}(2+i)} &= \frac{(-3+i)(2-i)}{\sqrt{2}(2+i)(2-i)} = \frac{-6+3i+2i-1}{(\sqrt{2})5} \\ &= \frac{-5+5i}{\sqrt{25}} = \frac{-1}{\sqrt{2}} + i\frac{1}{\sqrt{2}} \end{aligned}$$



# Solution

So  $\left| \frac{-3+i}{\sqrt{2}(2+i)} \right| = \left| \frac{-1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right| = \sqrt{(-1/\sqrt{2})^2 + (1/\sqrt{2})^2} = 1$  and  
 $\text{Arg} \left( \frac{-3+i}{\sqrt{2}(2+i)} \right) = \text{Arg} \left( \frac{-1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) = \frac{3\pi}{4}$ . It follows that the  
solutions of  $z^3 = \frac{-3+i}{\sqrt{2}(2+i)}$  are

$$z_0 = \cos(\pi/4) + i \sin(\pi/4) = \frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}}$$

$$z_1 = z_0 u_1 = \left( \frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) \left( -\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)$$

$$= \frac{-1 - \sqrt{3}}{2\sqrt{2}} + i \frac{\sqrt{3} - 1}{2\sqrt{2}}$$



# Solution

and

$$\begin{aligned} z_2 &= z_0 u_2 = \left( \frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) \left( -\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) \\ &= \frac{-1 + \sqrt{3}}{2\sqrt{2}} + i \frac{-\sqrt{3} - 1}{2\sqrt{2}}. \end{aligned}$$






# Solution

and

$$\begin{aligned} z_2 &= z_0 u_2 = \left( \frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) \left( -\frac{1}{2} - i \frac{\sqrt{3}}{2} \right) \\ &= \frac{-1 + \sqrt{3}}{2\sqrt{2}} + i \frac{-\sqrt{3} - 1}{2\sqrt{2}}. \end{aligned}$$



# Problem 1 from the exam from June 2009

Find all complex numbers  $z = x + iy$  which satisfy the equality  $|z + 1 - i\sqrt{3}| = |z - 1 + i\sqrt{3}|$ . Draw the solutions in a diagram.




# Solution



NTNU  
Norwegian University of  
Science and Technology

# Solution

Let  $w = 1 - i\sqrt{3}$ .




# Solution

Let  $w = 1 - i\sqrt{3}$ . Then

$|z + 1 - i\sqrt{3}| = |z - (-w)|$  is the distance between  $z$  and  $-w$ , and

$|z - w| = |z - 1 + i\sqrt{3}|$  is the distance between  $z$  and  $w$ .




# Solution

Let  $w = 1 - i\sqrt{3}$ . Then

$|z + 1 - i\sqrt{3}| = |z - (-w)|$  is the distance between  $z$  and  $-w$ , and

$|z - w| = |z - 1 + i\sqrt{3}|$  is the distance between  $z$  and  $w$ .

So  $|z + 1 - i\sqrt{3}| = |z - 1 + i\sqrt{3}|$  if and only if  $z$  has the same distance to  $-w$  as it has to  $w$ .



# Solution

Let  $w = 1 - i\sqrt{3}$ . Then


$|z + 1 - i\sqrt{3}| = |z - (-w)|$  is the distance between  $z$  and  $-w$ , and

$|z - w| = |z - 1 + i\sqrt{3}|$  is the distance between  $z$  and  $w$ .

So  $|z + 1 - i\sqrt{3}| = |z - 1 + i\sqrt{3}|$  if and only if  $z$  has the same distance to  $-w$  as it has to  $w$ .


The set of complex numbers  $z$  which satisfy the equality

$|z + 1 - i\sqrt{3}| = |z - 1 + i\sqrt{3}|$  is therefore the set of points that lie on the line  $L$  which goes through 0 and which is perpendicular to the line between  $w$  and  $-w$ .



# Solution


If we rotate the point  $w$  by 90 degrees, then we get a point that lies on the line  $L$ .





# Solution


If we rotate the point  $w$  by 90 degrees, then we get a point that lies on the line  $L$ . Since rotating  $w$  by 90 degrees is the same as multiplying  $w$  by  $i$ , it follows that  $wi = \sqrt{3} + i$  lies on the line  $L$ .



# Solution

If we rotate the point  $w$  by 90 degrees, then we get a point that lies on the line  $L$ . Since rotating  $w$  by 90 degrees is the same as multiplying  $w$  by  $i$ , it follows that  $wi = \sqrt{3} + i$  lies on the line  $L$ .

So the points that lie on the line  $L$  are the points of the form  $t(\sqrt{3} + i)$  where  $t$  is a real number.




# Solution

If we rotate the point  $w$  by 90 degrees, then we get a point that lies on the line  $L$ . Since rotating  $w$  by 90 degrees is the same as multiplying  $w$  by  $i$ , it follows that  $wi = \sqrt{3} + i$  lies on the line  $L$ .

So the points that lie on the line  $L$  are the points of the form  $t(\sqrt{3} + i)$  where  $t$  is a real number.

Thus,  $|z + 1 - i\sqrt{3}| = |z - 1 + i\sqrt{3}|$  if and only if  $z = t(\sqrt{3} + i)$  for some real number  $t$ .



# Next week's lectures



NTNU  
Norwegian University of  
Science and Technology

# Next week's lectures

Wednesday we shall



NTNU  
Norwegian University of  
Science and Technology

# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,



NTNU  
Norwegian University of  
Science and Technology

# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,
- look at *the fundamental theorem of algebra*,



NTNU  
Norwegian University of  
Science and Technology

# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,
- look at *the fundamental theorem of algebra*,
- introduce *the complex exponential function*,



NTNU  
Norwegian University of  
Science and Technology



# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,
- look at *the fundamental theorem of algebra*,
- introduce *the complex exponential function*,
- and study extensions of trigonometric functions to the complex numbers.



NTNU  
Norwegian University of  
Science and Technology

# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,
- look at *the fundamental theorem of algebra*,
- introduce *the complex exponential function*,
- and study extensions of trigonometric functions to the complex numbers.

Thursday we shall



NTNU  
Norwegian University of  
Science and Technology

# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,
- look at *the fundamental theorem of algebra*,
- introduce *the complex exponential function*,
- and study extensions of trigonometric functions to the complex numbers.

Thursday we shall

- study second-order differential equations,



# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,
- look at *the fundamental theorem of algebra*,
- introduce *the complex exponential function*,
- and study extensions of trigonometric functions to the complex numbers.

Thursday we shall

- study second-order differential equations,
- introduce the *Wronskian*,



# Next week's lectures

Wednesday we shall

- use complex numbers to solve polynomial equations,
- look at *the fundamental theorem of algebra*,
- introduce *the complex exponential function*,
- and study extensions of trigonometric functions to the complex numbers.

Thursday we shall

- study second-order differential equations,
- introduce the *Wronskian*,
- completely solve second-order homogeneous linear differential equations with constant coefficients.

