

# 11.7-8 Fourier Transform

Eugenia Malinnikova, NTNU

September 22, 2016

# Fourier Transform and Inverse Fourier Transform

Let  $f$  be a piece-wise continuous function with finite integral  $\int_{-\infty}^{\infty} |f(x)| dx$ , we define the Fourier transform of  $f$  by

$$\mathcal{F}(f)(w) = \hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ixw} dx$$

It is a continuous function. If  $\hat{f}$  is also has finite integral  $\int_{-\infty}^{\infty} |\hat{f}(w)| dw < \infty$  then, at each point  $x$  where  $f$  is continuous,

$$f(x) = \mathcal{F}^{-1}(\hat{f})(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{f}(w) e^{ixw} dw.$$

This is called the inversion formula. It allows to reconstruct the function from its Fourier transform and gives a representation of a function as a combination of the exponential ones.

- ▶ The Fourier transform is linear  $\mathcal{F}(af + bg) = a\mathcal{F}(f) + b\mathcal{F}(g)$ .
- ▶ The Fourier transform of the derivatives  
 $\mathcal{F}(f')(w) = iw\mathcal{F}(f)(w)$     $\mathcal{F}(f'')(w) = -w^2\mathcal{F}(f)(w)$
- ▶ Time shift  $\mathcal{F}(f(x - a))(w) = e^{-iaw}\mathcal{F}(f)(w)$
- ▶ Frequency shift  $\mathcal{F}(f)(w - b) = \mathcal{F}(f(x)e^{ibx})(w)$
- ▶ Convolution  $\mathcal{F}(f * g) = \sqrt{2\pi}\mathcal{F}(f)\mathcal{F}(g)$ ,

$$f * g(x) = \int_{-\infty}^{\infty} f(y)g(x - y)dy$$

# Example 1

## Example

Let  $f(x) = 1$ ,  $-1 < x < 1$  and  $f(x) = 0$  otherwise (see also the last lecture). Then

$$\hat{f}(w) = \frac{i}{\sqrt{2\pi}w} (e^{-iw} - e^{iw}) = \sqrt{\frac{2}{\pi}} \frac{\sin w}{w}$$

Then the inversion formula gives

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{f}(w) e^{ixw} dw = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\sin w}{w} e^{ixw} dw$$

Separating the real and imaginary parts, we obtain

$$\int_{-\infty}^{\infty} \frac{\sin w \cos xw}{w} = \begin{cases} \pi, & -1 < x < 1 \\ 1/2, & x = \pm 1 \\ 0, & |x| > 1 \end{cases}$$

## Example 2

### Example

Let  $f(x) = e^{-|x|}$  then

$$\begin{aligned}\hat{f}(w) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-|x|-iwx} dx = \\ &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-|x|} \cos xw dx + \frac{i}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-|x|} \sin xw dx\end{aligned}$$

Since  $f(x)$  is an even function, the second integral equals zero and we have

$$\hat{f}(w) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} e^{-|x|} \cos wx dx = \hat{f}_c(w)$$

the last integral is called the cosine Fourier transform (the calculation works for even function  $f$ ). Note that  $\hat{f}_c$  is even by the definition.

## Example 2, continued

We compute the Fourier transform

$$\hat{f}(w) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \frac{e^{-(1-iw)x} + e^{-(1+iw)x}}{2} dx = \sqrt{\frac{2}{\pi}} \frac{1}{1+w^2}$$

The inversion formula then implies

$$e^{-|x|} = f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \hat{f}_c(w) \cos wx dw = \frac{2}{\pi} \int_0^{\infty} \frac{\cos wx}{1+w^2} dx$$

# Real form of the Fourier transform

For general  $f$  we write its Fourier transform  $\hat{f}(w)$  as

$$\frac{1}{\sqrt{2\pi}} \left( \int_{-\infty}^{\infty} f(x) \cos wx dx - i \int_{-\infty}^{\infty} f(x) \sin wx dx \right) = a(w) - ib(w)$$

where  $a$  is even and  $b$  is odd. (Notation in the book:

$A = \sqrt{2/\pi}a, B = \sqrt{2\pi}b$ .) Then the inversion formula gives

$$\begin{aligned} f(x) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (a(w) - ib(w)) e^{iwx} dw = \\ &= \sqrt{\frac{2}{\pi}} \int_0^{\infty} a(w) \cos wx dw + \sqrt{\frac{2}{\pi}} \int_0^{\infty} b(w) \sin wx dw \end{aligned}$$

# Fourier transform of even and odd functions

For even function, we have seen that

$$\hat{f}(w) = \hat{f}_c(w) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \cos wx \, dx$$

and the function is reconstructed from its Fourier Cosine transform as

$$f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \hat{f}_c(w) \cos wx \, dx$$

Similarly, if  $f$  is odd, we have

$$\hat{f}(w) = -i\hat{f}_s(w), \quad \hat{f}_s(w) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} f(x) \sin wx \, dx$$

and we use the Sine Fourier transform for reconstruction

$$f(x) = \sqrt{\frac{2}{\pi}} \int_0^{\infty} \hat{f}_s(w) \sin wx \, dx$$



## One more example

Compute the Fourier transform of  $f(x) = e^{-x^2/2}$ . Let  $g(w) = \mathcal{F}(f)(w)$  then

$$\begin{aligned}g'(w) &= \frac{i}{\sqrt{2\pi}} \int_{-\infty}^{\infty} (-xe^{-x^2/2})e^{-iwx} dx = \\ &= i\mathcal{F}(f')(w) = i(-iw)\mathcal{F}(f)(w) = -wg(w)\end{aligned}$$

Then  $g'(w) = -wg(w)$  and  $g(w) = Ce^{-w^2/2}$ . From the inversion formula, we obtain  $C^2 = 1$  and it is not difficult to see that  $C > 0$  then  $C = 1$ .

More generally,

$$\mathcal{F}(e^{-ax^2}) = \frac{1}{\sqrt{2a}} e^{-w^2/(4a)}$$

# Applications of the Fourier transform

- ▶ Signal analysis, filters
- ▶ Differential equations and PDE
- ▶ Convolution equations