



Norwegian University of
Science and Technology

Department of Mathematical Sciences

Examination paper for **TMA4120 Calculus 4K**

Academic contact during examination: Eugenia Malinnikova, Yurii Lyubarskii

Phone: eugenia@math.ntnu.no, yura@math.ntnu.no

Examination date: one day in November

Examination time (from–to): 4 hours

Permitted examination support material: You need nothing but a pen/pencil, your head and a good mood! All the formulas you may need are in the attachment, see the last pages of the exam paper. However a simple calculator and Rottmann are still allowed on the final exam.

Other information:

This examination paper contains six problems with ten parts all together. Approximately a half of them is on Laplace transform, Fourier series and transform and their applications to differential equations and half on complex analysis. The problem parts are counted equally, though to pass the exam you should better pass both halves. Good luck!

Language: English

Number of pages: 1

Number of pages enclosed: 2

Checked by:

Date

Signature

Problem 1 Solve the initial value problem

$$y'' + 4y' + 4y = u(t - 1), \quad y(0) = 0, \quad y'(0) = 1.$$

Problem 2 We consider the following boundary value problem

$$u_{tt} + 2u_t + u = c^2 u_{xx}, \quad 0 < x < L, \quad t > 0, \quad u(0, t) = u(L, t) = 0. \quad (*)$$

- a) Find all solutions of (*) of the form $u(x, t) = F(x)G(t)$.
- b) Compute the coefficients of the sine Fourier series $S_f(x) = \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L}$ of the function $f(x)$ defined by $f(x) = x(L - x)$ for $0 < x < L$.
- c) Find a solution of (*) that satisfies the initial conditions

$$u(x, 0) = f(x), \quad u_t(x, 0) = 0,$$

where f is a function defined in b).

Problem 3 Show that if $\hat{f}(w)$ is the Fourier transform of a function $f(x)$ then the Fourier transform of the function $f(x) \sin bx$ is equal to $\frac{i}{2}(\hat{f}(w + b) - \hat{f}(w - b))$.

Compute the Fourier transform of the function $g(x) = \begin{cases} \sin 2x, & |x| \leq 1 \\ 0, & |x| > 1 \end{cases}$

Problem 4 Let $z_0 = i$. Determine the real and imaginary parts of

$$1 + z_0; \quad \frac{1 + z_0}{1 - z_0}; \quad \operatorname{Ln} z_0; \quad \operatorname{Ln}(1 + z_0).$$

Problem 5 Let $f(z) = \frac{\cos z}{z^2 + 1}$.

- a) Find all zeros and singular points of $f(z)$, classify the singularities.
- b) Compute $\oint_C f(z) dz$, where C is the circle of radius 3 centered at $z_0 = 1$.
- c) Let $g(w) = f(1/w)$ show that $g(w)$ has essential singularity at zero. (Hint: consider the values of g on a set $0 < |w| < r$.)

Problem 6 Evaluate the integral $\int_0^{\infty} \frac{\cos 3x}{x^4 + 3x^2 + 2} dx$.

Miscellaneous

- **Heaviside function** $u(t) = \begin{cases} 1, & t \geq 0 \\ 0, & t < 0 \end{cases}$, $u(t-a) = \begin{cases} 1, & t \geq a \\ 0, & t < a \end{cases}$
- **Dirac Delta function** $\delta(t-a)$ is zero everywhere except a and satisfies $\int_{-\infty}^{\infty} \delta(t-a)dt = 1$, moreover $\int_{-\infty}^{\infty} g(t)\delta(t-a) = g(a)$ for any continuous function g .
- **Convolution** For functions defined on the real line:
 $f * g(x) = \int_{-\infty}^{\infty} f(y)g(x-y)dy = \int_{-\infty}^{\infty} f(x-y)g(y)dy$, $-\infty < x < \infty$;
 for functions defined only on the positive half-axis:
 $f * g(x) = \int_0^x f(y)g(x-y)dy$.

Laplace transform

- $\mathcal{L}\{f\}(s) = F(s) \int_0^{\infty} f(t)e^{-st}dt$
- $\mathcal{L}\{e^{at}f(t)\}(s) = F(s-a)$
- $\mathcal{L}\{f'\}(s) = s\mathcal{L}\{f\}(s) - f(0)$
- $\mathcal{L}\{f''\}(s) = s^2\mathcal{L}\{f\}(s) - sf(0) - f'(0)$
- $\mathcal{L}\left\{\int_0^t f(\tau)d\tau\right\}(s) = \frac{1}{s}\mathcal{L}\{f\}(s)$
- $\mathcal{L}\{f * g\} = \mathcal{L}\{f\}\mathcal{L}\{g\}$
- $\mathcal{L}\{f(t-c)u(t-c)\} = e^{-cs}F(s)$,
 $c > 0$
- $\mathcal{L}\{tf(t)\}(s) = -F'(s)$
- $\mathcal{L}\left\{\frac{f(t)}{t}\right\}(s) = \int_s^{\infty} F(\sigma)d\sigma$

| $f(t)$ | $F(s)$ |
|-------------------------------|---------------------------|
| 1 | $\frac{1}{s}$ |
| $t^n, n = 1, 2, \dots$ | $\frac{n!}{s^{n+1}}$ |
| e^{at} | $\frac{1}{s-a}$ |
| $t^n e^{at}, n = 1, 2, \dots$ | $\frac{n!}{(s-a)^{n+1}}$ |
| $\cos bt$ | $\frac{s}{s^2+b^2}$ |
| $\sin bt$ | $\frac{b}{s^2+b^2}$ |
| $e^{at} \cos bt$ | $\frac{s-a}{(s-a)^2+b^2}$ |
| $e^{at} \sin bt$ | $\frac{b}{(s-a)^2+b^2}$ |
| $u(t-c), c > 0$ | $\frac{e^{-cs}}{s}$ |
| $\delta(t-c), c > 0$ | e^{-cs} |

Fourier series and Fourier transform

- Periodic functions with period $2L$, real and complex form

$$f(x) \sim a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi}{L}x + b_n \sin \frac{n\pi}{L}x \right) \sim \sum_{n=-\infty}^{\infty} c_n e^{in\pi x/L}$$

$$a_0 = \frac{1}{2L} \int_{-L}^L f(x) dx, \quad a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi}{L}x dx, \quad b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi}{L}x dx$$

$$c_n = \frac{1}{2L} \int_{-L}^L f(x) e^{-in\pi x/L} dx$$

- Parseval's identities

$$\frac{1}{2L} \int_{-L}^L |f(x)|^2 dx = \sum_{n=-\infty}^{\infty} |c_n|^2, \quad \int_{-\infty}^{\infty} |f(x)|^2 dx = \int_{-\infty}^{\infty} |\hat{f}(w)|^2 dw$$

- $\hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-iwx} dx$

- $f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \hat{f}(w) e^{iwx} dw$

- $\widehat{f'}(w) = iw \hat{f}(w)$

- $\widehat{f''}(w) = -w^2 \hat{f}(w)$

- $\widehat{f(x-a)}(w) = e^{-iaw} \hat{f}(w)$

- $\hat{f}(w-b) = e^{ibw} \widehat{f(x)}(w)$

- $\widehat{f * g} = \sqrt{2\pi} \hat{f} \hat{g}$

| $f(x)$ | $\hat{f}(w)$ |
|---|--|
| $\delta(x-a)$ | $\frac{1}{\sqrt{2\pi}} e^{-iaw}$ |
| $\begin{cases} 1, & x \leq a \\ 0, & x > a \end{cases}$ | $\sqrt{\frac{2}{\pi}} \frac{\sin aw}{w}$ |
| $\begin{cases} e^{-ax}, & x \geq 0 \\ 0, & x < 0 \end{cases}$ | $\frac{1}{\sqrt{2\pi}(a+iw)}$ |
| $\frac{1}{x^2+a^2}$ | $\sqrt{\frac{\pi}{2}} \frac{e^{-a w }}{a}$ |
| e^{-ax^2} | $\frac{1}{\sqrt{2a}} e^{-w^2/(4a)}$ |

Complex numbers and analytic functions

- $e^{x+iy} = e^x (\cos y + i \sin y)$,

$$\cos z = \frac{e^{iz} + e^{-iz}}{2}, \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}, \quad \cosh z = \frac{e^z + e^{-z}}{2}, \quad \sinh z = \frac{e^z - e^{-z}}{2}$$

- Taylor and Laurant series of an analytic function

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n, \quad a_n = \frac{f^{(n)}(z_0)}{n!} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^{n+1}} dz$$

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n + \sum_{n=1}^{\infty} \frac{b_n}{(z - z_0)^n}, \quad b_n = \frac{1}{2\pi i} \oint_C f(z) (z - z_0)^{n-1} dz$$