



Norwegian University of
Science and Technology

Department of Mathematical Sciences

Examination paper for **TMA4120 Calculus 4K**

Academic contact during examination:

Phone:

Examination date: 11.08.2018

Examination time (from–to): 9:00-13:00

Permitted examination support material:

Language: English

Number of pages: 5

Number of pages enclosed: 3

Checked by:

Informasjon om trykking av eksamensoppgave	
Originalen er:	
1-sidig <input type="checkbox"/>	2-sidig <input checked="" type="checkbox"/>
sort/hvit <input checked="" type="checkbox"/>	farger <input type="checkbox"/>
skal ha flervalgskjema <input type="checkbox"/>	

Date

Signature

Problem 1 Let

$$g(t) = \begin{cases} 1, & 0 \leq t < 1 \\ -1, & 1 \leq t < 2 \\ 0, & 2 < t. \end{cases}$$

a) Find the Laplace transform of g .

Løsning By definition:

$$\mathcal{L}(g) = \int_0^1 e^{-st} dt - \int_1^2 e^{-st} dt = \left(-\frac{1}{s} e^{-st} \right) \Big|_0^1 + \left(-\frac{1}{s} e^{-st} \right) \Big|_1^2.$$

Thus $\mathcal{L}(g) = s^{-1}(1 - 2e^{-s} - e^{-2s})$.

Alternatively, $g(t) = u(t) - 2u(t-1) + u(t-2)$, using the tabel, we obtain

$$\mathcal{L}(g) = \mathcal{L}(u(t)) - 2\mathcal{L}(u(t-1)) + \mathcal{L}(u(t-2)) = \frac{1 - 2e^{-s} + e^{-2s}}{s}.$$

b) Solve the initial value problem

$$y''(t) + 4y(t) = g(t), \quad t \geq 0, \quad y(0) = 0, \quad y'(0) = 0.$$

Løsning We apply the Laplace transform, let $Y = \mathcal{L}(y)$, then using the initial data and computation from part a), we obtain

$$(s^2 + 4)Y = \frac{1 - 2e^{-s} + e^{-2s}}{s}, \quad Y(s) = \frac{1 - 2e^{-s} + e^{-2s}}{s(s^2 + 4)}.$$

We use partial fraction decomposition,

$$Y(s) = (1 - 2e^{-s} + e^{-2s}) \left(\frac{1}{4s} - \frac{s}{s^2 + 4} \right).$$

Then we apply the inverse Laplace transform

$$\underline{y(t) = \frac{1 + \cos 2t}{4}u(t) - \frac{1 + \cos 2(t-1)}{2}u(t-1) + \frac{1 + \cos 2(t-2)}{4}u(t-2).}$$

Problem 2 Consider the boundary value problem:

$$\begin{cases} u_t = u_{xx}, & t > 0, \ 0 < x < 1 \\ u(0, t) = u(1, t) = 0 \end{cases}$$

- a) Find all solutions on the form $u(x, t) = F(t)G(x)$ satisfying the end point conditions $G(0) = G(1) = 0$.

Løsning We divide the variables in the equation:

$$\frac{F'}{F} = \frac{G''}{G} = k.$$

Taking into account the boundary condition, we want to find $G(x)$ such that $G'' = kG$ and $G(0) = G(1) = 0$. We know that it is possible only when $k < 0$ and $k = -p^2$. Then $G(x) = \sin px$ and the condition $G(1) = 0$ implies $p = \pi n$ for a positive integer n .

Then $F' = -(\pi n)^2 F$ and $F(t) = Ce^{-(\pi n)^2 t}$. all solutions with divided variables are of the form

$$\underline{u_n(x, t) = C_n \sin(\pi n x) e^{-(\pi n)^2 t}.$$

- b) Use the principle of superposition to find the solution of the boundary value problem that also satisfies the following initial condition

$$u(x, 0) = 3 \sin(\pi x) + 5 \sin(4\pi x), \quad 0 < x < 1.$$

Løsning We know that a linear combination of solutions is a solution (superposition principle), to satisfy the the initial condition we take

$$\underline{u(x, t) = 3 \sin(\pi x) e^{-\pi^2 t} + 5 \sin(4\pi x) e^{-(4\pi)^2 t}.$$

Problem 3 Calculate the Fourier transform of the function

$$f(x) = \begin{cases} e^{-2018x}, & x \geq 0 \\ 0, & x < 0. \end{cases}$$

Løsning By the definition

$$\hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_0^\infty e^{-2018x - ixw} dx = \frac{1}{\sqrt{2\pi}} \frac{1}{2018 + iw} = \frac{1}{\sqrt{2\pi}} \frac{2018 - iw}{2018^2 + w^2}.$$

Problem 4 Assume that $f(z)$ is an analytic function in a domain Ω that satisfies $|f(z)| = 1$ for all $z \in \Omega$. Prove that f is a constant.

Løsning Consider the function $h(z) = \ln f(z)$, since f has no zeros in Ω h is defined and analytic in Ω . Let $h(z) = u(z) + iv(z)$, where u and v are the real and imaginary parts of h . We have

$$e^{u(z)}e^{iv(z)} = f(z).$$

Taking the absolute values we get $e^{u(z)} = |f(z)| = 1$. It implies that $u(z) = 0$ for all $z \in \Omega$. Then from the Cauchy-Riemann equation we see that $v_x = v_y = 0$ and v is also a constant. Finally we conclude that $f(z) = e^{h(z)}$ is a constant.