

TMA4165: PROBLEM SHEET II

published: 28/01/2021

1. (June 2018 examination) Let  $x^0(t)$  and  $x^\eta(t)$  be solutions to the Cauchy problem

$$0 = \dot{x} - x + \frac{1}{1+r^2}x^2, \quad x(0) = \frac{1}{2},$$

with  $r = 0$  and  $r = \eta > 0$ , respectively.

Show that for each  $r \in \mathbb{R}$ , this equation has a unique solution on some interval of existence around  $t = 0$ , and that the dependence on the parameter  $r$  is continuous, in particular, that there is a bounded function  $K : t \rightarrow \mathbb{R}$  independent of  $r$  such that

$$|x^0(t) - x^\eta(t)| \leq K(t)\eta^2.$$

2. Show that

$$T(x) = \frac{\pi}{2} + x - \arctan(x), \quad x \in \mathbb{R}$$

has no fixed point even though

$$|T(x) - T(y)| < |x - y|$$

for every pair of distinct  $x, y \in \mathbb{R}$ .

Explain if this contradicts the contraction mapping principle.

3. Let  $f : [0, 1] \rightarrow \mathbb{R}$  be a continuous function. Show that the following *boundary value* problem for a *second-order* ODE:

$$\begin{aligned} -\frac{d^2}{dx^2}u(x) + \lambda \sin(u(x)) &= f(x), & x \in [0, 1] \\ u(0) &= 0, & u(1) = 0 \end{aligned}$$

has a unique solution for every sufficiently small  $\lambda$ . Write out the first three Picard iterations, beginning with  $u_0 \equiv 0$ .

4. (*Perko* Example 2.1) Show that the Cauchy problem

$$\frac{d}{dt}y(t) = y^2(t) - t, \quad y(0) = 1$$

exhibits finite-time blow-up, and that this maximal time of existence is less than  $t = 1.23$ .

5. The following equation for  $f : [-a, a] \rightarrow \mathbb{R}$  arises in a model for one-dimensional gas dynamics:

$$f(x) = 1 + \frac{1}{\pi} \int_{-a}^a \frac{1}{1 + (x - y)^2} f(y) dy, \quad x \in [-a, a].$$

Show that the integral equation has a unique, non-negative solution for every  $0 < a < \infty$ . What happens in the asymptotic regime as  $a \rightarrow \infty$ ?

6. Exercise 7 in Chapter 4 of *Schaeffer and Cain*, or prove the following, more general version:

Suppose  $f : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{R}^d$  is Lipschitz in its first argument and uniformly continuous in its second with modulus of continuity  $\omega$ . For solutions  $\mathbf{x}^\alpha(t)$  and  $\mathbf{y}^\beta(t)$  on the interval  $[0, T]$ , with initial conditions  $\mathbf{x}^\alpha(0) = \mathbf{x}_0$  and  $\mathbf{y}^\beta(0) = \mathbf{y}_0$ , the following bound holds:

$$|\mathbf{x}_\alpha(t) - \mathbf{y}_\beta(t)| \leq C_{T,f} |\mathbf{x}_0 - \mathbf{y}_0| \omega(|\alpha - \beta|),$$

where  $C_{T,f}$  is a constant depending on  $T$  and  $f$ .