

## WEEK 9

This exercise set is on eigentheory of the Fourier transform (see page 173 in [3]). For notations, see the second part of TMA4170 lecture notes.

**Exercise 1:** We have already proved that the Hermite functions  $h_k$  are orthogonal to each other and satisfy

$$\sum_{k=0}^{\infty} h_k(x) \frac{t^k}{k!} = e^{-\pi(x+2t)^2} \cdot e^{2\pi t^2}.$$

Use this formula to prove that

$$\|h_k\|^2 := \int_{\mathbb{R}} |h_k(x)|^2 dx = 2^{-1/2} (4\pi)^k k!.$$

**Exercise 2:** Use integration by parts to show that

$$((A - B)h_{k-1}, (A - B)h_{k-1}) = -(A + B)(A - B)h_{k-1}, h_{k-1}),$$

i.e. the adjoint operator of  $A - B$  is  $-(A + B)$ .

**Exercise 3:** Recall that  $h_k = (A - B)h_{k-1}$  and

$$(A + B)(A - B) = 4\pi f - 2\pi.$$

Use Exercise 2 and

$$f(h_{k-1}) = -(k - \frac{1}{2})h_{k-1}$$

to prove that

$$\|h_k\|^2 = 4\pi k \|h_{k-1}\|^2.$$

Then use it to evaluate  $\|h_k\|^2$  a second way.

**Remark:** Notice that  $f(h_k) = -(k + \frac{1}{2})h_k$  is equivalent to

$$-(h_k)''(x) + 4\pi^2 x^2 h_k(x) = 2\pi(2k + 1)h_k(x).$$

Consider

$$e_k(x) = \frac{h_k((2\pi)^{-1/2}x)}{\|h_k((2\pi)^{-1/2}x)\|} = \frac{h_k((2\pi)^{-1/2}x)}{\pi^{1/4} \sqrt{(4\pi)^k k!}},$$

we know that  $\{e_k\}$  is an orthonormal basis of  $L^2(\mathbb{R})$ . Try to do the following two exercises.

**Exercise 4:** Put  $Lf = -f'' + x^2 f$ . Show that

$$L(e_k) = (2k + 1)e_k, \quad k = 0, 1, 2, \dots$$

**Exercise 5:** One version of the Heisenberg uncertainty principle (see Week 8 Exercise 2) is:

$$(Lf, f) = \|f'\|^2 + \|xf\|^2 \geq \|f\|^2, \quad \forall f \in \mathcal{S}.$$

Try to prove the following effective version

$$(Lf, f) = \|f'\|^2 + \|xf\|^2 = \sum_{n=0}^{\infty} (2n+1) |(f, e_n)|^2 \geq \sum_{n=0}^{\infty} |(f, e_n)|^2 = \|f\|^2,$$

for all  $f \in \mathcal{S}$ . In particular, the above formula gives

$$\|f'\|^2 + \|xf\|^2 \geq \|f\|^2 + 2 \cdot \text{Distance}\{f, \mathbb{C}e_0\}^2,$$

where  $\text{Distance}\{f, \mathbb{C}e_0\}$  denotes the  $L^2$  distance from  $f$  to the complex line  $\mathbb{C}e_0$  generated by the Gaussian function  $e_0$ .

**Remark:** For more information on the above Hermite operator  $L$  and the related representation theory, see [2]. For an effective version of the original Heisenberg uncertainty principle, see page 117 in [1].

#### REFERENCES

- [1] H. Dym and H. P. McKean, *Fourier Series and Integrals*, Academic Press, 1972.
- [2] R. Howe, *On the role of the Heisenberg group in harmonic analysis*, Bull. Amer. Math. Soc. (1980).
- [3] E. Stein and R. Shakarchi, *Fourier analysis*, Princeton lectures in analysis.