



1 (*Properties of the lower limit*)

a) Let $(y_k)_{k \in \mathbb{N}}, (z_k)_{k \in \mathbb{N}} \subset \mathbb{R}$ be two real sequences. Show that

$$\liminf_{k \rightarrow \infty} y_k + \liminf_{k \rightarrow \infty} z_k \leq \liminf_{k \rightarrow \infty} (y_k + z_k).$$

In addition, find an example where this inequality is strict.

b) Let I be any index set (possibly infinite), and let $(y_k^i)_{k \in \mathbb{N}} \subset \mathbb{R}, i \in I$, be a family of sequences. Show that

$$\sup_{i \in I} \liminf_{k \rightarrow \infty} y_k^i \leq \liminf_{k \rightarrow \infty} (\sup_{i \in I} y_k^i).$$

2 Let I be any index set and let $f_i: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ be lower semi-continuous. Show that the function $f: \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ given as

$$f(x) = \sup_{i \in I} f_i(x)$$

is lower semi-continuous.

3 For the following functions, decide whether they are lower semi-continuous or coercive, and whether they attain a global minimizer:

a) The function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = x^4 - 20x^3 + \sup_{k \in \mathbb{N}} \sin(kx).$$

b) The function $g: \mathbb{R} \rightarrow \mathbb{R}$ given by

$$g(x) = e^x - \frac{1}{x^2 + 1}.$$

c) The function $h: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$h(x) = x_1^2(1 + x_2^3) + x_1^2.$$

4 For which constants $c \in \mathbb{R}$ does the function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) := (x^2 - 2x + c)e^{-x^2}$$

attain a global minimum?

- 5 Given a matrix $A \in \mathbb{R}^{n \times n}$, we denote by

$$\|A\|_F := \left(\sum_{i,j=1}^n a_{ij}^2 \right)^{1/2}$$

its *Frobenius norm*. Show that the optimisation problem

$$\min_{\substack{A \in \mathbb{R}^{n \times n} \\ \det A > 0}} \left(\|A\|_F + \frac{1}{\det A} \right)$$

admits a global minimum.

- 6 (See *N&W, Exercise 2.1*). The *Rosenbrock function* is defined as

$$f(x) := 100(x_2 - x_1^2)^2 + (1 - x_1)^2.$$

- a) Compute the gradient and the Hessian of the Rosenbrock function.
- b) Show that the point $(1, 1)$ is the unique (global and local) minimizer of f .