



1 Consider the function

$$f(x, y, z) = 2x^2 + xy + y^2 + yz + z^2 - 6x - 7y - 8z + 9.$$

- a) Using the first order necessary conditions, find a critical point of f .
- b) Use the second-order sufficient conditions in order to verify that this point is a local minimum.
- c) Prove that this point is a global minimum of f .

2 Assume that f is a continuously differentiable function satisfying

$$\lim_{\|x\| \rightarrow \infty} \frac{f(x)}{\|x\|} = +\infty.$$

Show that the equation

$$\nabla f(x) = u$$

has a solution for every $u \in \mathbb{R}^n$.

Hint: Consider global minima of the function $f_u(x) := f(x) - u^T x$.

3 Show that the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f(x, y) = \log(e^x + e^y)$$

is convex.

4 (Arithmetic–Geometric Mean Inequality) Let $\alpha, \beta > 0$ be such that $\alpha + \beta = 1$. Show that for all $x, y > 0$ one has

$$x^\alpha y^\beta \leq \alpha x + \beta y$$

with equality if and only if $x = y$.

Hint: show that the function $-\log x$ is strictly convex on $\mathbb{R}_{>0}$.

5 (See *N&W, Exercise 2.8*) Assume that $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is a convex function. Show that the set of minimisers of f is convex.

6 Show that a strictly convex function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ has at most one global minimiser. In addition, find a strictly convex function that has no global minimiser at all.

7 Assume that $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is a strictly convex and continuous function that attains a minimiser. Show that f is coercive.¹

Hint: assuming x^ is the minimiser of f , show that $f(x) \geq c\|x - x^*\|$ whenever $\|x - x^*\| \geq 1$, where $c > 0$ is the minimum of f on $\{x : \|x - x^*\| = 1\}$.*

¹The continuity of f is strictly speaking not a necessary requirement, as it can be shown that every convex function defined on the whole space \mathbb{R}^n is necessarily continuous.