Exercise 6

February 20, 2024

Problem 1.

Sketch the region

$$\Omega = \left\{(x, y) \in \mathbb{R}^2 : y \geq x \text{ and } y^4 \leq x^3\right\}$$

and compute the tangent cone and the set of linearised feasible directions for each point in Ω. For which points in Ω is the LICQ satisfied?

Solution.

We first define the constraint functions,

$$c_1(x, y) = y - x \quad \text{and} \quad c_2(x, y) = x^3 - y^4,$$

so that $\Omega = \{(x, y) \in \mathbb{R}^2 : c_1(x, y) \geq 0 \text{ and } c_2(x, y) \geq 0\}$, and sketch of the region Ω is displayed in the Figure below.



Figure 1: Region Ω in grey, with colors on the boundary specifiying the active constraints.

In order to characterise the tangent cone $T_\Omega(x, y)$ and the set of linearised feasible directions $\mathcal{F}(x, y)$, we employ Lemma 12.2 in N&W, which states that if the LICQ condition holds at a feasible point (x, y), then $T_\Omega(x, y) = \mathcal{F}(x, y)$.

Note first that the LICQ condition holds vacuously in the interior of Ω because all constraints are inactive, and therefore, $T_\Omega(x,y) = \mathcal{F}(x,y) = \mathbb{R}^2$ at interior points.

Next we consider boundary points with precisely one active constraint. Starting with points for which $c_1(x,y) = 0$, and excluding $(0,0)$ and $(1,1)$ where also c_2 is active. We find that $\nabla c_1(x,y) = (-1,1)$. Since $\nabla c_1 \neq 0$, the LICQ condition holds, and so

$$\begin{aligned} T_\Omega(x,y) = \mathcal{F}(x,y) &= \left\{d = (d_1,d_2) \in \mathbb{R}^2 : \nabla c_1(x,y)^\top d \geq 0\right\} \\ &= \left\{d = (d_1,d_2) \in \mathbb{R}^2 : d_2 \geq d_1\right\}. \end{aligned}$$

Similarly, if only c_2 is active, we observe that the LICQ condition holds because $\nabla c_2(x,y) = (3x^2, -4y^3) \neq 0$ away from $(0,0)$. This yields

$$\begin{aligned} T_\Omega(x,y) = \mathcal{F}(x,y) &= \left\{d = (d_1,d_2) \in \mathbb{R}^2 : \nabla c_2(x,y)^\top d \geq 0\right\} \\ &= \left\{d = (d_1,d_2) \in \mathbb{R}^2 : 3x^2 d_1 \geq 4y^3 d_2\right\}. \end{aligned}$$

The constraint gradients at $(1,1)$ equal $\nabla c_1 = (-1,1)$ and $\nabla c_2 = (3,-4)$, which are linearly independent. Thus the LICQ condition is true, and

$$\begin{aligned} T_\Omega(1,1) = \mathcal{F}(1,1) &= \left\{d = (d_1,d_2) \in \mathbb{R}^2 : \nabla c_1(1,1)^\top d \geq 0 \text{ and } \nabla c_2(1,1)^\top d \geq 0\right\} \\ &= \left\{d = (d_1,d_2) \in \mathbb{R}^2 : 3d_1 \geq 4d_2\right\}. \text{(I think it should be} \{2d_1 \geq 3d_2\}) \end{aligned}$$

Lastly, since $\nabla c_1(0,0) = (-1,1)$ and $\nabla c_2(0,0) = 0$, the LICQ condition fails at $(0,0)$, and we cannot expect that $T_\Omega(0,0) = \mathcal{F}(0,0)$. Readily,

$$\begin{aligned} \mathcal{F}(0,0) &= \left\{d \in \mathbb{R}^2 : \nabla c_1(0,0)^\top d \geq 0 \text{ and } \nabla c_2(0,0)^\top d \geq 0\right\} \\ &= \left\{d \in \mathbb{R}^2 : d_2 \geq d_1\right\}. \end{aligned}$$

In order to find the tangent cone, we first consider limiting directions along the constraint boundaries $c_1(x,y) = 0$ and $c_2(x,y) = 0$ as $(x,y) \to (0,0)$. Travelling towards $(0,0)$ when c_1 is active, we may put, using the notation in N&W,

$$z_k = (1/k, 1/k) \quad \text{and} \quad t_k = 1/k,$$

and obtain the limiting direction

$$d = \lim_{k\to\infty} \frac{z_k - (0,0)}{t_k} = (1,1).$$

Note: the length of d is irrelevant; we only care about its direction. Similarly, travelling along $c_2(x,y) = 0$ yields $d = (0,1)$, using for example, the sequences

$$z_k = \left(k^{-1/3}, k^{-1/4}\right) \quad \text{and} \quad t_k = k^{-1/4}.$$

It can furthermore be seen that approaching $(0,0)$ from the interior of Ω gives tangent directions "between" these borderline cases, and so

$$T_\Omega(0,0) = \{d \in \mathbb{R}^2 : d_2 \geq d_1 \geq 0\}.$$

Problem 2.

Assume that one wants to solve the optimisation problem

$$\max_x f(x) \qquad \text{such that} \qquad \begin{cases} c_i(x) = 0 & \text{for all } i \in \mathcal{E}, \\ c_i(x) \geq 0 & \text{for all } i \in \mathcal{I}. \end{cases}$$

How can we modify the KKT conditions such that one obtains (first order) necessary conditions for this maximisation problem?

Solution.

Let

$$\mathcal{L}(x,\lambda) = f(x) - \sum_{i \in \mathcal{E} \cup \mathcal{I}} \lambda_i c_i(x)$$

be the Lagrangian associated with the maximisation problem. Since solving $\max_x f(x)$ is equivalent to solving $\min_x -f(x)$, we can state the KKT conditions for the minimisation problem. To this end, let

$$\widehat{\mathcal{L}}(x,\mu) = -f(x) - \sum_{i \in \mathcal{E} \cup \mathcal{I}} \mu_i c_i(x)$$

be the Lagrangian for the minimisation problem, so that the KKT conditions become

$$\begin{aligned} -\nabla f(x) - \sum_{i \in \mathcal{E} \cup \mathcal{I}} \mu_i \nabla c_i(x) = \nabla_x \widehat{\mathcal{L}}(x,\mu) = 0, & \\ c_i(x) = 0 \quad \text{for all } i \in \mathcal{E}, & \\ c_i(x) \geq 0 \quad \text{for all } i \in \mathcal{I}, & \\ \mu_i \geq 0 \quad \text{for all } i \in \mathcal{I}, & \\ \mu_i c_i(x) = 0 \quad \text{for all } i \in \mathcal{E} \cup \mathcal{I}. & \end{aligned}$$

Since

$$\mathcal{L}(x,-\mu) = -\widehat{\mathcal{L}}(x,\mu) \quad \text{and} \quad \nabla_x \mathcal{L}(x,-\mu) = -\nabla_x \widehat{\mathcal{L}}(x,\mu),$$

we see that changing the signs of the Lagrange multipliers, that is, putting $\lambda = -\mu$, is the only modification in the KKT conditions for the maximisation problem.

Problem 3.

Consider the constrained optimization problem

$$\min_{(x,y)}(x^2+y^2) \qquad \text{such that} \qquad \begin{cases} x+y \geq 1, \\ y \leq 2, \\ y^2 \geq x. \end{cases}$$

a) Formulate the KKT-conditions for this optimisation problem.

b) Find all KKT points.

c) Find all local and global minima.

(Part b) can be very tedious. One strategy is to consider all possible active sets and determine for each active set whether KKT-points exist. It can also be extremely helpful to sketch the feasible set and the function in order to identify the different possibilities for active sets.)

Solution.

a) We begin by stating the problem in standard form, writing $\mathbf{x} = [x, y]^T$:

$$\min_{\mathbf{x}\in\mathbb{R}^2} f(\mathbf{x}) \quad \text{s.t.} \quad c_i(\mathbf{x}) \geq 0, \quad i = 1, 2, 3,$$

where

$$\begin{aligned} f(\mathbf{x}) &= x^2 + y^2, \\ c_1(\mathbf{x}) &= x + y - 1, \\ c_2(\mathbf{x}) &= 2 - y, \\ c_3(\mathbf{x}) &= y^2 - x. \end{aligned}$$

The KKT conditions can now be stated as follows:

$$2x^* - \lambda_1^* + \lambda_3^* = 0 \tag{1a}$$
$$2y^* - \lambda_1^* + \lambda_2^* - 2y^*\lambda_3^* = 0 \tag{1b}$$
$$x^* + y^* - 1 \geq 0 \tag{1c}$$
$$2 - y^* \geq 0 \tag{1d}$$
$$y^{*2} - x^* \geq 0 \tag{1e}$$
$$\lambda_i^* \geq 0, \quad i = 1, 2, 3 \tag{1f}$$
$$\lambda_1^*(x^* + y^* - 1) = 0 \tag{1g}$$
$$\lambda_2^*(2 - y^*) = 0 \tag{1h}$$
$$\lambda_3^*(y^{*2} - x^*) = 0. \tag{1i}$$



Figure 2: Feasible set. Note: The lower "triangle" extends further toward infinity.

b) The feasible set is sketched in Figure 2.

We will find all KKT points by systematically considering all possible active sets of constraints. Remember that a constraint c_i is active at a point $\mathbf{x}$ if $c_i(\mathbf{x}) = 0$. Also, the LICQ conditions are satisfied at every point we consider here; with one active constraint, the LICQ conditions hold trivially, and in the cases with two constraints it is not hard to check that the LICQ conditions do hold.

Observe that if $\mathbf{x}^* = [x^*, y^*]^T$ is a KKT point, then from (1a) and (1b) we have:

$$x^* = \frac{\lambda_1^* - \lambda_3^*}{2}, \quad y^* = \frac{\lambda_1^* - \lambda_2^*}{2(1-\lambda_3^*)}.$$

From here on, we will drop the asterisk in the notation and write x for x^*, etc.

First, suppose that the active set is empty, i.e. neither of (1c)-(1e) are equalities. This corresponds to the interior of the domain. Then, by (1g)-(1i), we have $\lambda_1 = \lambda_2 = \lambda_3 = 0$, and so $x = y = 0$. But this point is not feasible, since it violates condition (1c). Thus, with the active set empty, there are no KKT points.

Next, we consider the case when the active set contains one index, i.e. exactly one of (1c)-(1e) is an equality. This corresponds to the boundaries of the domain, excepting the corner points. If (1c) is ac-

tive, then $\lambda_2 = \lambda_3 = 0$ while $\lambda_1 \geq 0$. We get

$$x = \frac{\lambda_1}{2}, \quad y = \frac{\lambda_1}{2},$$

and inserting this into (1c) (which is now an equality), we get the condition

$$\frac{\lambda_1}{2} + \frac{\lambda_1}{2} - 1 = 0 \Rightarrow \lambda_1 = 1,$$

giving us the point $(x, y) = (\frac{1}{2}, \frac{1}{2})$. But this point violates condition (1e), so $(\frac{1}{2}, \frac{1}{2})$ is not a KKT point.

If (1d) is active, then $\lambda_1 = \lambda_3 = 0$ while $\lambda_2 \geq 0$, so

$$x = 0, \quad y = -\frac{\lambda_2}{2}.$$

Inserting this into the equality (1d), we get

$$2 + \frac{\lambda_2}{2} = 0 \Rightarrow \lambda_2 = -4.$$

Since the Lagrange multiplier is negative, KKT conditions are not satisfied at this point.

If (1e) is active, then $\lambda_1 = \lambda_2 = 0$ while $\lambda_3 \geq 0$, so

$$x = -\frac{\lambda_3}{2}, \quad y = 0.$$

Inserting this into the equality (1e), we get

$$\frac{\lambda_3}{2} = 0 \Rightarrow \lambda_3 = 0.$$

This gives the candidate point $(0, 0)$, which is not feasible since it violates (1c), and thereby is not a KKT point.

Having considered all possible active sets of one index, we now turn to the cases with two indices, i.e. exactly two of (1c)-(1e) are equalities. This corresponds to the corner points of the domain. First, if (1c) and (1d) are both active, then $\lambda_3 = 0$ while $\lambda_1, \lambda_2 \geq 0$. This gives us

$$x = \frac{\lambda_1}{2}, \quad y = \frac{\lambda_1 - \lambda_2}{2}.$$

Plugging this into equalities (1c) and (1d) yields:

$$\begin{aligned} \frac{\lambda_1}{2} + \frac{\lambda_1 - \lambda_2}{2} - 1 &= 0 \\ 2 - \frac{\lambda_1 - \lambda_2}{2} &= 0, \end{aligned}$$

with solutions $\lambda_1 = -2$ and $\lambda_2 = -6$. Since the multipliers are negative, this is not a KKT point.

Next, if (1c) and (1e) are both active, then $\lambda_2 = 0$ while $\lambda_1, \lambda_3 \geq 0$, which means

$$x = \frac{\lambda_1 - \lambda_3}{2}, \quad y = \frac{\lambda_1}{2(1-\lambda_3)}.$$

Plugging this into equalities (1c) and (1e) yields:

$$\frac{\lambda_1 - \lambda_3}{2} + \frac{\lambda_1}{2(1-\lambda_3)} - 1 = 0$$
$$\frac{\lambda_1^2}{4(1-\lambda_3)^2} - \frac{\lambda_1 - \lambda_3}{2} = 0.$$

Solving this set of equations yields $\lambda_1 = 5 \pm \frac{9}{\sqrt{5}}$ and $\lambda_3 = 2 \pm \frac{4}{\sqrt{5}}$, thereby giving the candidate points $(x, y) = (\frac{1}{2}(3 \pm \sqrt{5}), \frac{1}{2}(-1 \mp \sqrt{5}))$ which both satisfy the KKT conditions. Since $\lambda_1, \lambda_3 \geq 0$, these points are minimizer candidates. Note: This result can be arrived upon by the easier approach of first finding the points (x, y) where c_1 and c_3 are both active, then working out what λ_1 and λ_3 are.

Finally, we check the case where (1d) and (1e) are both active, i.e. $\lambda_1 = 0$ while $\lambda_2, \lambda_3 \geq 0$. This gives us

$$x = -\frac{\lambda_3}{2}, \quad y = -\frac{\lambda_2}{2(1-\lambda_3)}.$$

Plugging this into equalities (1d) and (1e) yields:

$$2 + \frac{\lambda_2}{2(1-\lambda_3)} = 0$$
$$\frac{\lambda_2^2}{4(1-\lambda_3)^2} + \frac{\lambda_3}{2} = 0,$$

which can be solved to find $\lambda_2 = -28$ and $\lambda_3 = -8$. Since the multipliers are negative, this is not a KKT point.

Concerning the case with all constraints active, we may conclude that no KKT point exists; all three constraint functions cannot be active at the same point. The investigation is summarized in the table below.

Point	λ_1	λ_2	λ_3	KKT?
(0,2)	0	-4	0	No
$(\frac{1}{2}(3+\sqrt{5}), \frac{1}{2}(-1-\sqrt{5}))$	$5 + \frac{9}{\sqrt{5}}$	0	$2 + \frac{4}{\sqrt{5}}$	Yes
$(\frac{1}{2}(3-\sqrt{5}), \frac{1}{2}(-1+\sqrt{5}))$	$5 - \frac{9}{\sqrt{5}}$	0	$2 - \frac{4}{\sqrt{5}}$	Yes
(-1,2)	-2	-6	0	No
(4,2)	0	-28	-8	No

c) To determine whether the KKT points that are minimizer candidates are in fact local minimizers, we check the second order sufficient conditions from Theorem 12.6 in N&W, i.e. whether

$$w^T \nabla^2_{xx} \mathcal{L}(x, \lambda) w > 0 \ \forall w \in \mathcal{C}(x, \lambda), w \neq 0, \tag{2}$$

where, $\mathcal{C}(x, \lambda)$ is the critical cone at x, given by (12.53) in N&W.

For both candidates, i.e. $(\frac{1}{2}(3 \pm \sqrt{5}), \frac{1}{2}(-1 \mp \sqrt{5}))$, we have that the critical cone is simply given as $\mathcal{C}(x, \lambda) = \{0\}$. This is because any $w \in \mathcal{C}(x, \lambda)$ must be orthogonal to the $\nabla c_i(x)$ for which $\lambda_i > 0$, of which there are two for each point. Since the LICQ conditions hold at both points, these two vectors are linearily independent and thus span $\mathbb{R}^2$. The only vector orthogonal to $\mathbb{R}^2$ is the zero vector. Thereby, the only vector in $\mathcal{C}(x, \lambda)$ is the zero vector for these points, and thus condition (2) holds. We can conclude that $(\frac{1}{2}(3 \pm \sqrt{5}), \frac{1}{2}(-1 \mp \sqrt{5}))$ are strict local minimizers.

We note that $f(\frac{1}{2}(3 - \sqrt{5}), \frac{1}{2}(-1 + \sqrt{5})) < f(\frac{1}{2}(3 + \sqrt{5}), \frac{1}{2}(-1 - \sqrt{5}))$ and $f(\mathbf{x}) \to \infty$ in the unbounded region of the feasible domain. This means that $(\frac{1}{2}(3 - \sqrt{5}), \frac{1}{2}(-1 + \sqrt{5}))$ is a global minimizer and $(\frac{1}{2}(3 + \sqrt{5}), \frac{1}{2}(-1 - \sqrt{5}))$ is a local minimizer.

Problem 4.

Consider the constrained optimization problem

$$\min_{(x,y)} x \qquad \text{such that} \qquad \begin{cases} y \geq x^4, \\ y \leq x^3. \end{cases}$$

Find all KKT points and local minima for this optimisation problem.

Solution.

We begin by stating the problem in standard form, writing $\mathbf{x} = [x, y]^T$:

$$\min_{\mathbf{x} \in \mathbb{R}^2} f(\mathbf{x}) \quad \text{s.t.} \quad c_i(\mathbf{x}) \geq 0, \quad i = 1, 2,$$

where

$$\begin{aligned} f(\mathbf{x}) &= x, \\ c_1(\mathbf{x}) &= y - x^4 \\ c_2(\mathbf{x}) &= x^3 - y. \end{aligned}$$

The KKT conditions for this problem can be stated as follows:

$$
\begin{aligned}
1 + 4x^3\lambda_1 - 3x^2\lambda_2 &= 0, && \text{(3a)}\\
-\lambda_1 + \lambda_2 &= 0, && \text{(3b)}\\
y - x^4 &\geq 0, && \text{(3c)}\\
x^3 - y &\geq 0, && \text{(3d)}\\
\lambda_i &\geq 0, \quad i = 1, 2, && \text{(3e)}\\
\lambda_1(y - x^4) &= 0, && \text{(3f)}\\
\lambda_2(x^3 - y) &= 0.. && \text{(3g)}
\end{aligned}
$$

Now, we can take a shortcut; from (3b), we see that $\lambda_1 = \lambda_2$, and from (3a) we see that there cannot exist any KKT point for which $\lambda_1 = \lambda_2 = 0$. Therefore, the cases with no active constraints ($\lambda_1 = \lambda_2 = 0$) and one active constraint ($\lambda_1 = 0$ or $\lambda_2 = 0$) cannot produce KKT points. We are left with considering the case where both constraints are active, i.e. the corner points (0,0) and (1,1).

In the point (1,1), we find (by (3a) and (3b)) that $\lambda_1 = \lambda_2 = -1$, and therefore this is not a KKT point.

The last point is $(0, 0)$, for which we cannot write the gradient of f at $(0, 0)$ (which is $[1, 0]^T$) as a non-negative linear combination of the gradients of the constraints, and which therefore is not a KKT point (here we can simply write $(0, 0)$ does not satisfy the condition 3a, therefore, $(0, 0)$ cannot be a KKT point). This does not, however, mean that it is not a minimizer. Applying common sense, it is clearly a local minimum, as no other points with $x = 0$ are feasible, and $x = 0$ is the lowest possible value of the objective function.

Problem 5.

Consider the constrained optimization problem

$$
\min_{(x,y)} \left(-x^2 - (y-1)^2\right) \qquad \text{such that} \qquad \begin{cases} y \geq Cx^2, \\ y \leq 2, \end{cases}
$$

where $C > 0$ is some positive parameter.

a) Show that the point $(0, 0)$ is a KKT point for all parameters $C > 0$ and that the LICQ is satisfied at $(0, 0)$.

b) Formulate the second order necessary and sufficient optimality conditions for the point $(0, 0)$. For which parameters C are these conditions satisfied? For which parameters C is the point $(0, 0)$ a local minimum?

Solution.

a) Introducing

$$f(x,y) = -x^2-(y-1)^2, \quad c_1(x,y) = y-Cx^2, \qquad \text{and} \quad c_2(x,y) = 2-y,$$

the minimisation problem becomes

$$\min_{x,y} f(x,y) \qquad \text{subject to} \qquad c_1(x,y) \geq 0 \text{ and } c_2(x,y) \geq 0.$$

Let also

$$\mathcal{L}(x,y,\lambda_1,\lambda_2) = f(x,y) - \lambda_1 c_1(x,y) - \lambda_2 c_2(x,y)$$

be the Lagrangian, with multipliers λ_1 and λ_2.

Focusing on the KKT conditions, it is clear that $(0,0)$ is feasible. Moreover, from the complementarity conditions

$$\lambda_1 c_1(0,0) = 0 \qquad \text{and} \qquad \lambda_2 c_2(0,0) = 0,$$

we require $\lambda_2 = 0$ because c_2 is inactive. (Note: c_1 is active, so the first condition holds.) Computing

$$\nabla_{x,y}\mathcal{L}(x,y,\lambda_1,0) = \nabla f(x,y) - \lambda_1 \nabla c_1(x,y) = \begin{bmatrix} -2x(1-\lambda_1 C) \\ -2(y-1) - \lambda_1 \end{bmatrix}$$

and demanding that this gradient vanishes at $(x,y) = (0,0)$, then give $\lambda_1 = 2$, with no restriction on C. Hence, $(0,0)$ is a KKT point for all $C > 0$. Additionally, since c_1 is the only active constraint and $\nabla c_1(0,0) = (0,1) \neq 0$, it follows that the LICQ is satisfied as well.

b) Let $\mathcal{C}$ be the critical cone at $(0,0)$ with Lagrange multipliers $(\lambda_1,\lambda_2) = (2,0)$. Then $(0,0)$, which satisfies the KKT conditions, is a local minimiser of the constrained problem only if (*necessary condition*) the Lagrangian Hessian

$$\nabla^2_{(x,y)}\mathcal{L}(0,0,2,0) = \begin{bmatrix} -2(1-2C) & 0 \\ 0 & -2 \end{bmatrix}$$

is positive semi-definite on $\mathcal{C}$, that is,

$$w^\top \nabla^2_{(x,y)}\mathcal{L}(0,0,2,0) w \geq 0 \quad \text{for all} \quad w \in \mathcal{C}.$$

If (*sufficient condition*), however, this Hessian is positive definite on $\mathcal{C}$, then $(0,0)$ is a (strict) local minimiser.

Since c_1 is the only active constraint and $\lambda_1 > 0$, we find that

$$\mathcal{C} = \{(w_1, w_2) \in \mathbb{R}^2 : \nabla c_1(0,0)^\top w = 0\} = \{(w_1, 0) \in \mathbb{R}^2 : w_1 \in \mathbb{R}\}.$$

Therefore, with $w = (w_1, 0) \in \mathcal{C}$,

$$w^\top \nabla^2_{(x,y)} \mathcal{L}(0,0,2,0) w = -2(1-2C)w_1^2,$$

which is nonnegative if and only if $C \geq 1/2$, and strictly positive for all $w \in \mathcal{C} \setminus \{0\}$ if and only if $C > 1/2$. Thus $(0,0)$ is a (strict) local minimum whenever $C > 1/2$, but cannot be a minimiser if $0 < C < 1/2$. It remains to examine $C = 1/2$. To this end, we consider, for example, points (x, y) approaching $(0,0)$ along $c_1(x,y) = 0$, that is, points for which $y = x^2/2 \to 0$. This yields

$$f\left(x, \tfrac{1}{2}x^2\right) = -x^2 - \left(\tfrac{1}{2}x^2 - 1\right)^2 = -\tfrac{1}{4}x^4 - 1,$$

which is strictly less than $f(0,0) = -1$ for all $x \neq 0$. In particular, $(0,0)$ is not a local minimiser when $C = 1/2$.