

## Trial exam

Spring 2020

1 Let  $X = \{a, b, c, d\}$ .

Which of the following collections of subsets of  $X$  is *not* a topology on  $X$ ?

Select one alternative:

- $\mathcal{T} = \{\emptyset, \{a\}, \{a, b\}, \{a, b, d\}, \{a, d\}, \{d\}, X\}$
- $\mathcal{T} = \{\emptyset, \{a\}, \{a, b, c\}, \{a, d\}, \{b, c, d\}, X\}$
- $\mathcal{T} = \{\emptyset, \{b, c, d\}, \{d\}, X\}$
- $\mathcal{T} = \{\emptyset, \{a\}, \{a, b\}, \{a, c, d\}, X\}$

**Solution:** The collection

$$\mathcal{T} = \{\emptyset, \{a\}, \{a, b, c\}, \{a, d\}, \{b, c, d\}, X\}$$

is *not* a topology on  $X$  as T3 is not satisfied: the intersection of the open sets  $\{a, b, c\}$  and  $\{b, c, d\}$  is not an open set as  $\{a, b, c\} \cap \{b, c, d\} = \{b, c\} \notin \mathcal{T}$ . (Similarly,  $\{a, d\} \cap \{b, c, d\} \notin \mathcal{T}$ .)

2 Let  $X = \{a, b, c, d, e\}$  and let  $\mathcal{T} = \{\emptyset, \{a\}, \{a, b, c\}, \{b, c\}, X\}$  be a topology on  $X$ .

Which of the following statements is *false*?

Select one alternative:

- The interior of  $\{d, e\}$ ,  $\text{Int}(\{d, e\})$ , is equal to the empty set  $\emptyset$ .
- The closure of  $\{e\}$ ,  $\overline{\{e\}}$ , is equal to  $\{d, e\}$ .
- $X$  is compact.
- $X$  is Hausdorff.

**Solution:** Since the points  $b$  and  $c$  are not separated by disjoint open sets,  $X$  is *not* Hausdorff. (The same argument applies to  $a$  and  $d$ ,  $a$  and  $e$ ,  $b$  and  $d$ ,  $b$  and  $e$ ,  $c$  and  $d$ ,  $c$  and  $e$ , and  $d$  and  $e$ .)

3 Let  $X$  be a topological space, and let  $\mathcal{B}$  be a basis for the topology on  $X$ . Furthermore, let  $A$  be a subset of  $X$ .

Show that  $x \in \overline{A}$  if and only if  $B \cap A \neq \emptyset$  for every basis element  $B \in \mathcal{B}$  where  $x \in B$ .

(Here  $\overline{A}$  denotes the closure of  $A$ .)

**Solution:** This is essentially the same problem as Exercise 3.6 in [lecture notes](#). We first show that  $x \notin \overline{A}$  if and only if there is no neighborhood  $U$  of  $x$  such that  $U \cap A = \emptyset$ .

Let  $x \notin \overline{A}$ . Then  $x \in \overline{A}^c = X \setminus \overline{A}$ . Since  $\overline{A}^c$  is open in  $X$ ,  $\overline{A}^c$  is a neighborhood of  $x$  such that  $\overline{A}^c \cap A = \emptyset$ .

For the other implication assume that there is a neighborhood  $U$  of  $x$  such that  $U \cap A = \emptyset$ . Since  $U$  is open in  $X$ ,  $U^c$  is closed in  $X$ , and  $A \subseteq U^c$ . Thus as the closure of  $A$  is the smallest closed subset of  $X$  that contains  $A$ , it follows that  $\overline{A} \subset U^c$ . Hence,  $x \notin \overline{A}$ .

From what we have just proved it follows that  $x \notin \overline{A}$  if and only if there is a neighborhood  $U$  of  $x$  such that  $U \cap A = \emptyset$ . Since each basis element  $B \in \mathcal{B}$  is open in  $X$ , and for each neighborhood  $U$  of  $x$  there is a basis element  $B \in \mathcal{B}$  such that  $x \in B \subseteq U$ , we have  $x \in \overline{A}$  if and only if there is a basis element  $B \in \mathcal{B}$  such that  $B \cap A \neq \emptyset$  and  $x \in B$ .

- 4 Let  $\mathbb{R}$  be the set of real numbers equipped with the standard topology, and consider the set of rational numbers  $\mathbb{Q}$  as a subspace of  $\mathbb{R}$ .

Show that the subset  $A = \{x \in \mathbb{Q} \mid -\sqrt{5} < x < \sqrt{5}\}$  of  $\mathbb{Q}$  is both open and closed in  $\mathbb{Q}$ .

**Solution:** Since the open interval  $(-\sqrt{5}, \sqrt{5})$  is open in  $\mathbb{R}$  and  $A = (-\sqrt{5}, \sqrt{5}) \cap \mathbb{Q}$ , it follows that  $A$  is open in the subspace topology on  $\mathbb{Q}$ . Similarly, since the closed interval  $[-\sqrt{5}, \sqrt{5}]$  is closed in  $\mathbb{R}$  and  $A = [-\sqrt{5}, \sqrt{5}] \cap \mathbb{Q}$ , it follows that  $A$  is closed in the subspace topology on  $\mathbb{Q}$ .

- 5 Let  $X = \{a, b, c, d\}$  be given the topology  $\mathcal{T}_X = \{\emptyset, \{a\}, \{a, c, d\}, \{c, d\}, X\}$ , and let  $Y = \{1, 2, 3\}$  be given the topology  $\mathcal{T}_Y = \{\emptyset, \{1\}, \{1, 3\}, Y\}$ .

Find a basis for the product topology on  $X \times Y$  expressed using bases for the topologies on  $X$  and  $Y$ , respectively.

**Solution:** Note that  $\mathcal{B}_X = \{\{a\}, \{c, d\}, X\}$  is a basis for  $\mathcal{T}_X$ , and that  $\mathcal{B}_Y = \{\{1\}, \{1, 3\}, Y\}$  is a basis for  $\mathcal{T}_Y$ . Then by Theorem 5.8 in the [lecture notes](#), it follows that

$$\mathcal{B}_{X \times Y} = \{B_X \times B_Y \mid B_X \in \mathcal{B}_X \text{ and } B_Y \in \mathcal{B}_Y\}$$

is a basis for the product topology on  $X \times Y$ .

- 6 Let  $X$  be a topological space, and consider  $I = [0, 1]$  as a subspace of  $\mathbb{R}$  where  $\mathbb{R}$  is given the standard topology. Furthermore, let the *cone on  $X$*  be the quotient space  $CX = X \times I / \sim$ , where  $\sim$  is the equivalence relation on the product space  $X \times I$  given by  $(x, 0) \sim (x', 0)$  for all  $x, x' \in X$ .

Show that  $CX$  is path connected.

**Solution:** Denote by  $[x, t]$  the equivalence class of  $(x, t)$ , and let  $v$  denote the vertex of  $CX$ , i.e.,  $v$  is the point corresponding to  $[X, 0]$ . Then for any  $x \in X$  and any  $t > 0$ , there is a path  $f: I \rightarrow X \times I$  given by

$$f(s) = (x, (1 - s)t).$$

In other words,  $f$  is a path in  $X \times I$  from  $(x, t)$  to  $(x, 0)$ . Thus  $\pi \circ f: I \rightarrow CX$  is a path in  $CX$  from  $[x, t]$  to  $v$ . Hence,  $CX$  is path connected.

- 7 Let  $X$  be a topological space, and let  $A_1, A_2, \dots, A_n$  be subspaces of  $X$  each of which is compact in  $X$ .

Show that  $\bigcup_{i=1}^n A_i$  is compact in  $X$ .

**Solution:** Let  $\mathcal{A}$  be an open cover of  $\bigcup_{i=1}^n A_i$ , i.e.,

$$\bigcup_{i=1}^n A_i \subseteq \bigcup_{U \in \mathcal{A}} U.$$

We must show that  $\mathcal{A}$  has a finite subcover.

Note that  $A_i \subseteq \bigcup_{U \in \mathcal{A}} U$  for all  $i \in \{1, 2, \dots, n\}$ . Hence,  $\mathcal{A}$  is an open cover of each  $A_i$ . Since each  $A_i$  is compact in  $X$ , there is a finite subcover  $\mathcal{A}_i$  of  $\mathcal{A}$  that covers  $A_i$ . Thus

$$\bigcup_{i=1}^n \mathcal{A}_i$$

is a finite subcover of the open cover  $\mathcal{A}$ . Hence,  $\bigcup_{i=1}^n A_i$  is compact in  $X$ .

8 Let  $X$  be a contractible space.

Show that  $X$  is path connected.

**Solution:** Let  $a, b \in X$ . Since  $X$  is contractible there is a homotopy  $H: X \times I \rightarrow X$  such that

$$H(x, 0) = x$$

and

$$H(x, 1) = c$$

for all  $x \in X$  for some fixed point  $c \in X$ . Then the map  $\alpha: I \rightarrow X$  given by  $\alpha(t) = H(a, t)$  is a path in  $X$  from  $\alpha(0) = a$  to  $\alpha(1) = c$ . Similarly, the map  $\beta: I \rightarrow X$  given  $\beta(t) = H(b, t)$  is a path in  $X$  from  $\beta(0) = b$  to  $\beta(1) = c$ . Thus  $a, b$  and  $c$  are all joined by paths. Hence,  $X$  is path connected.

9 Let  $B$  be a Hausdorff space, and let  $E$  be a topological space.

Show that if  $p: E \rightarrow B$  is a covering map, then  $E$  must be Hausdorff.

**Solution:** Let  $e$  and  $e'$  be two distinct points in  $E$ , i.e.,  $e \neq e'$ . If  $p(e) \neq p(e')$ , there are, by assumption that  $B$  is Hausdorff, disjoint open subsets  $U$  and  $U'$  in  $X$  such that  $p(e) \in U$  and  $p(e') \in U'$ . Then  $p^{-1}(U)$  and  $p^{-1}(U')$  are disjoint neighborhoods of  $e$  and  $e'$ , respectively.

Let  $U$  be a neighborhood of  $p(e)$  that is evenly covered by  $p$ , i.e.,

$$p^{-1}(U) = \bigsqcup_{\lambda \in \Lambda} V_\lambda$$

where  $V_\lambda$  is open in  $E$ , and  $p|_{V_\lambda}: V_\lambda \rightarrow U$  is a homeomorphism for each  $\lambda \in \Lambda$ . If  $p(e) = p(e')$ , then there are distinct indices  $\lambda_1$  and  $\lambda_2$  in  $\Lambda$  such that  $e \in V_{\lambda_1}$  and  $e' \in V_{\lambda_2}$ .

Hence,  $E$  is Hausdorff.

10 Let  $n$  be an integer that is greater than or equal to 3, and let  $D^n = \{(x_1, x_2, \dots, x_n) \in \mathbb{R}^n \mid \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} \leq 1\}$  be considered as a subspace of  $\mathbb{R}^n$  where  $\mathbb{R}^n$  is given the standard topology.

Show that the inclusion map  $i: D^n \setminus \{0\} \rightarrow D^n$  induces an isomorphism of fundamental groups. (Here 0 denotes the origin in  $\mathbb{R}^n$ .)

You may assume as a known fact that the  $m$ -sphere  $S^m$  is simply connected where  $S^m$  is considered as a subspace of  $\mathbb{R}^{m+1}$  and  $m$  is an integer that is greater than or equal to 2.

**Solution:** Since  $D^n$  is contractible, and  $D^n \setminus \{0\}$  is a deformation retract of  $S^{n-1}$  (the homotopy  $H: D^n \setminus \{0\} \times I \rightarrow D^n \setminus \{0\}$  given by

$$H(x, t) = (1-t)x + t \frac{x}{\|x\|}, \quad x = (x_1, x_2, \dots, x_n), \quad \|x\| = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2},$$

is a deformation retraction of  $D^n \setminus \{0\}$  onto  $S^{n-1}$ ), it follows that for any  $d_0 \in D^n \setminus \{0\}$ , we have

$$\pi_1(D^n \setminus \{0\}, d_0) \cong 0$$

and

$$\pi_1(D^n, d_0) \cong 0$$

where 0 denotes the trivial group and where we have used the fact that  $S^{n-1}$  is simply connected. Hence, the induced homomorphism  $i_*: \pi_1(D^n \setminus \{0\}, d_0) \rightarrow \pi_1(D^n, d_0)$  is an isomorphism for each  $d_0 \in D^n \setminus \{0\}$ .