

Harmonicfunctionology

Harald Hanche-Olsen

The Laplace operator on \mathbb{R}^n

$$\Delta = \sum_{i=1}^n \frac{\partial^2}{\partial x_i^2}$$

plays a rôle in the wave and heat equations, but even more fundamentally, in the *Laplace equation*

$$\Delta u = 0 \tag{1}$$

and the *Poisson equation* $-\Delta u = f$ where f is a given function.

A C^2 solution of (1) is called *harmonic*. (Later, in Theorem 4, we will find that harmonic functions are in fact infinitely differentiable.)

Much is gained from the trivial observation that $\Delta u = \nabla \cdot \nabla u$ together with various applications of the divergence theorem or two of its corollaries, Green's first and second identities.

Let's get started by simply integrating Δu over a bounded domain ω with piecewise C^1 boundary:

$$\int_{\omega} \Delta u(\mathbf{x}) \, d^n \mathbf{x} = \int_{\omega} \nabla \cdot \nabla u(\mathbf{x}) \, d^n \mathbf{x} = \int_{\partial\omega} \partial_{\nu} u(\mathbf{x}) \, dS(\mathbf{x}). \tag{2}$$

This immediately proves

Proposition 1. *If a C^2 function u on a domain Ω is harmonic, then*

$$\int_{\partial\omega} \partial_{\nu} u \, dS = 0 \tag{3}$$

for all bounded domains ω with $\bar{\omega} \subset \Omega$ having piecewise C^1 boundary.

Conversely, if (3) holds for every ball $\omega = B(\mathbf{x}, r)$ whose closure lies within Ω , then u is harmonic.

Proof. We have already proved the first part. For the converse, (3) and (2) imply that the *average* of Δu over any ball is zero. By letting the radius of the ball $B(\mathbf{x}, r)$ tend to zero, we conclude that $\Delta u(\mathbf{x}) = 0$. ■

Definition. The (*radius r*) *spherical average* of a function u at a point \mathbf{x} is defined to be

$$\tilde{u}_{\mathbf{x}}(r) = \int_{\partial B(\mathbf{x}, r)} u \, dS = \int_{\mathbb{S}^{n-1}} u(\mathbf{x} + r\mathbf{y}) \, dS(\mathbf{y}),$$

where $\mathbb{S}^{n-1} \subset \mathbb{R}^n$ is the unit sphere and the “barred” integral signs denote the average:

$$\bar{\int}_{\partial B(\mathbf{x},r)} u \, dS = \frac{1}{A_n r^{n-1}} \int_{\partial B(\mathbf{x},r)} u \, dS,$$

where A_n is the area of \mathbb{S}^{n-1} . Note that the second integral in the definition of $\tilde{u}_{\mathbf{x}}(r)$ makes sense even for $r < 0$; thus, we adopt this as the definition for all real r for which the integrand is defined on \mathbb{S}^{n-1} . We see that $\tilde{u}_{\mathbf{x}}$ is an *even* function; it is C^k if u is C^k , and $\tilde{u}_{\mathbf{x}}(0) = u(\mathbf{x})$.

When ω is a ball, we can rewrite (2) in terms of spherical averages: Noting that the volume of the ball $B(\mathbf{x}, r)$ is $A_n r^n/n$, we find

$$\int_{B(\mathbf{x},r)} \Delta u(\mathbf{y}) \, d^n \mathbf{y} = \frac{n}{r} \int_{B(\mathbf{x},r)} \partial_{\nu} u(\mathbf{y}) \, dS(\mathbf{y}) = \frac{n}{r} \int_{\mathbb{S}^{n-1}} \partial_r u(\mathbf{x} + r\mathbf{y}) \, dS(\mathbf{y}),$$

where we can move the r derivative outside the integral, and arrive at

$$\int_{B(\mathbf{x},r)} \Delta u(\mathbf{y}) \, d^n \mathbf{y} = \frac{n}{r} \tilde{u}'_{\mathbf{x}}(r). \quad (4)$$

Along with $\tilde{u}_{\mathbf{x}}(0) = u(\mathbf{x})$, this implies

Theorem 2 (The mean value property of harmonic functions). *A C^2 function u on a domain Ω is harmonic if and only if $\tilde{u}_{\mathbf{x}}(r) = u(\mathbf{x})$ for all $\mathbf{x} \in \Omega$ and all r for which $\bar{B}(\mathbf{x}, |r|) \subset \Omega$.*

In general, we say a function u satisfies the *mean value property* if $\tilde{u}_{\mathbf{x}}(r) = u(\mathbf{x})$ whenever $\bar{B}(\mathbf{x}, |r|) \subset \Omega$. We shall see below (Theorem 4) that the mean value property characterizes harmonic functions. But first, we collect an easy consequence of (4).

Proposition 3. *For any C^2 function u , we have*

$$\Delta u(\mathbf{x}) = n \tilde{u}''_{\mathbf{x}}(0).$$

Proof. The function $\tilde{u}_{\mathbf{x}}$ is even, so $\tilde{u}'_{\mathbf{x}}(0) = 0$. Therefore, letting $r \rightarrow 0$ in (4), we arrive at the stated result. ■

Theorem 4 (The mean value property and regularity). *Assume that a continuous function u satisfies the mean value property on a domain Ω . Then u is infinitely differentiable, and is therefore harmonic. In particular, every harmonic function is infinitely differentiable.*

Proof. This proof may seem long, but only because we use it to develop some tools that have wider applicability.

First, define a *standard mollifier* $\rho: \mathbb{R}^n \rightarrow \mathbb{R}$. Here is one of many possible definitions:

$$\rho(\mathbf{x}) = \begin{cases} ae^{1/(|\mathbf{x}|^2-1)}, & |\mathbf{x}| < 1 \\ 0, & |\mathbf{x}| \geq 1, \end{cases}$$

where the constant $a > 0$ is chosen to ensure that

$$\int_{\mathbb{R}^n} \rho \, d\mathbf{x} = 1.$$

That is one of the defining qualities of a standard mollifier. The others are: That $\rho \geq 0$ everywhere, that it vanishes outside the unit ball, that it is infinitely differentiable, and is radially symmetric – that is, a function of $|\mathbf{x}|$ alone.

For any $\delta > 0$ we can squeeze the mollifier to fit inside a ball of radius δ :

$$\rho_\delta(\mathbf{x}) = \frac{1}{\delta^n} \rho\left(\frac{\mathbf{x}}{\delta}\right),$$

so that ρ_δ also has integral 1, but vanishes outside the ball $B(0, \delta)$.

Now we consider the convolution product

$$u * \rho_\delta(\mathbf{x}) = \int_{\mathbb{R}^n} u(\mathbf{y}) \rho_\delta(\mathbf{x} - \mathbf{y}) \, d^n \mathbf{y}.$$

This is defined for all $x \in \Omega$ with a distance less than δ to the complement of Ω . Thus, for any $x \in \Omega$, we can make δ small enough so that $u * \rho_\delta$ is defined at x .

Moreover, $u * \rho_\delta$ is infinitely differentiable: This is proved by differentiating with respect to the components of x under the integral sign, as much as you like.

Finally, the mean value property of u and the radial symmetry of ρ_δ combine to ensure that $u(\mathbf{x}) = u * \rho(\mathbf{x})$ for all x where $u * \rho$ is defined, which is what we were going to prove.

For a detailed argument, write

$$u * \rho(\mathbf{x}) = \int_{\mathbb{R}^n} u(\mathbf{x} - \mathbf{y}) \rho_\delta(\mathbf{y}) \, d^n \mathbf{y}$$

and write the integral in polar form:

$$\begin{aligned} u * \rho(\mathbf{x}) &= \int_0^\delta \int_{\partial B(\mathbf{x}, r)} u(\mathbf{x} - \mathbf{y}) \rho_\delta(\mathbf{y}) \, dS(\mathbf{y}) \, dr \\ &= \int_0^\delta \int_{\mathbb{S}^{n-1}} u(\mathbf{x} - r\mathbf{y}) \rho_\delta(r\mathbf{y}) \, dS(\mathbf{y}) \, r^{n-1} \, dr. \end{aligned} \tag{5}$$

Now use the radial symmetry: $\rho_\delta(r\mathbf{y})$ is constant for $\mathbf{y} \in \mathbb{S}^{n-1}$, so this factor can be moved outside the inner integral. Next, use the mean value property of u :

$$\int_{\mathbb{S}^{n-1}} u(\mathbf{x} - r\mathbf{y})\rho_\delta(r\mathbf{y}) d^n \mathbf{y} = A_n u(\mathbf{x}).$$

But $u(\mathbf{x})$ is a constant, which we move outside the outer integral in (5). We are left with

$$u * \rho(\mathbf{x}) = u(\mathbf{x}) \int_0^\delta r^{n-1} A_n \rho_\delta(r\mathbf{y}) dr$$

where \mathbf{y} is any unit vector. But running the whole calculation in reverse, this time without the u term, reveals that the integral here is merely the integral of ρ_δ , which has the value 1. Thus we are left with $u * \rho(\mathbf{x}) = u(\mathbf{x})$, as claimed. ■

The maximum principle

Definition. A C^2 function u is called *subharmonic* if $\Delta u \geq 0$, and *superharmonic* if $\Delta u \leq 0$. Thus it is harmonic if and only if it is both subharmonic and superharmonic. (The reason for the naming will become clear later; see Corollary 7.) Clearly, u is superharmonic if and only if $-u$ is subharmonic.

Theorem 5 (Strong maximum principle). *Assume that $u \in C^2(\Omega)$ is subharmonic in a region $\Omega \subseteq \mathbb{R}^n$. If u has a global maximum in Ω , then u is constant.*

Proof. Let M be the global maximum of u , and put

$$S = \{ \mathbf{x} \in \Omega \mid u(\mathbf{x}) = M \}.$$

Then S is a closed subset of Ω , by the continuity of u . It is also nonempty by assumption.

Consider any $\mathbf{x} \in S$. From (4) and the subharmonicity of u , we get $\tilde{u}'_{\mathbf{x}}(r) \geq 0$ for $r > 0$. Thus we get $\tilde{u}_{\mathbf{x}}(r) \geq \tilde{u}_{\mathbf{x}}(0) = u(\mathbf{x}) = M$ for $r > 0$ (so long as $\bar{B}(\mathbf{x}, r) \subset \Omega$). But $u \leq M$ everywhere, and if $u < M$ anywhere on the sphere $\partial B(\mathbf{x}, r)$, we would get $\tilde{u}_{\mathbf{x}}(r) < M$. Thus $u = M$ in some neighbourhood of \mathbf{x} . This means that S is open.

We have shown that S is an open, closed, and nonempty subset of the connected set Ω . Therefore $S = \Omega$, and the proof is complete. ■

Remark. Obviously, we obtain a strong *minimum* principle for *superharmonic* functions by multiplying by -1 . In particular, a non-constant harmonic function cannot attain a minimum or maximum value anywhere in Ω .

Corollary 6 (Weak maximum principle). *Let $\Omega \subset \mathbb{R}^n$ be a bounded domain, and assume that $u \in C(\overline{\Omega}) \cap C^2(\Omega)$ is subharmonic. Then*

$$\max \{ u(\mathbf{x}) \mid \mathbf{x} \in \overline{\Omega} \} = \max \{ u(\mathbf{x}) \mid \mathbf{x} \in \partial\Omega \}.$$

In particular, a harmonic function which is continuous on $\overline{\Omega}$ attains its minimum and maximum values on the boundary $\partial\Omega$.

Proof. The weak principle is an obvious consequence of the strong principle. However, it is worth noting that a much more elementary proof exists. Namely, for any $\varepsilon > 0$, let $v(\mathbf{x}) = u(\mathbf{x}) + \varepsilon|\mathbf{x}|^2$, and note that then $\Delta v > 0$. But $\Delta v(\mathbf{x}) \leq 0$ if \mathbf{x} is an interior minimum point for v , so v cannot have any maximum in the interior. Thus for any $\mathbf{x} \in \Omega$,

$$u(\mathbf{x}) = v(\mathbf{x}) - \varepsilon|\mathbf{x}|^2 \leq \max_{\partial\Omega} v \leq \max_{\partial\Omega} u + \varepsilon \max_{\mathbf{x} \in \partial\Omega} |\mathbf{x}|^2.$$

Now let $\varepsilon \rightarrow 0$ to arrive at the conclusion $u(\mathbf{x}) \leq \max_{\partial\Omega} u$. ■

Our next result explains the terms *sub-* and *super*harmonic: A subharmonic function is below, and a superharmonic above, a harmonic function given the same boundary data.

Corollary 7. *Assume that Ω is a bounded domain, that $u, v \in C(\overline{\Omega}) \cap C^2(\Omega)$, that u is harmonic in Ω , and that $v = u$ on $\partial\Omega$. If v is subharmonic, then $v \leq u$ in Ω , while if v is superharmonic, then $v \geq u$ in Ω .*

Proof. Apply the weak maximum principle to $v - u$ if v is subharmonic, or to $u - v$ if v is superharmonic. ■

Remark. Corollary 7 suggests a strategy for proving existence of a solution to the Dirichlet problem for the Laplace equation: Assume we are trying to find a harmonic function on Ω with the given boundary value g . Consider the pointwise *supremum* of all subharmonic functions which are $\leq g$ on $\partial\Omega$, and the pointwise *infimum* of all superharmonic functions which are $\geq g$ on $\partial\Omega$. If the two functions coincide, they should provide a solution to the problem. This is the basis for *Perron's method*, which we will hopefully get a look at later.

The Poisson equation

We now turn our study to the *Poisson equation*:

$$-\Delta u = f \tag{6}$$

where f is a known continuous function. (It *must* be continuous to allow for classical, i.e., C^2 , solutions u .)

Referring all the way back to (2), we quickly get the following generalization of Proposition 1:

Proposition 8. *A C^2 function u on a domain Ω solves the Poisson equation (6) if and only if*

$$-\int_{\partial\omega} \partial_\nu u(\mathbf{x}) \, dS(\mathbf{x}) = \int_{\omega} f(\mathbf{x}) \, d^n \mathbf{x} \tag{7}$$

for all bounded domains ω with $\bar{\omega} \subset \Omega$ having piecewise C^1 boundary. It is sufficient to consider balls $\omega = B(\mathbf{0}, r)$.

As an example, we consider a Poisson equation with a radially symmetric right hand side $f(\mathbf{x}) = \mathring{f}(|\mathbf{x}|)$. We expect to find a radially symmetric solution $u(\mathbf{x}) = \mathring{u}(|\mathbf{x}|)$. Now (7) with $\omega = B(\mathbf{0}, r)$ becomes

$$-A_n r^{n-1} \mathring{u}'(r) = A_n \int_0^r \mathring{f}(s) s^{n-1} \, ds.$$

Taking the derivative and rearranging turns this into the ODE

$$-\frac{1}{r^{n-1}} \frac{d}{dr} (r^{n-1} \mathring{u}'(r)) = \mathring{f}(r). \tag{8}$$

A direct calculation reveals that indeed,

$$\Delta \mathring{u}(|\mathbf{x}|) = \mathring{u}''(|\mathbf{x}|) + \frac{n-1}{|\mathbf{x}|} \mathring{u}'(|\mathbf{x}|) = \frac{1}{r^{n-1}} \frac{d}{dr} (r^{n-1} \mathring{u}'(r)) \Big|_{r=|\mathbf{x}|}$$

so a solution to (8) will in fact solve the Poisson equation in the radially symmetric case.

The ODE (8) is easily integrated:

$$-r^{n-1} \mathring{u}'(r) = \int_0^r \mathring{f}(s) s^{n-1} \, ds = \frac{1}{A_n} \int_{B(\mathbf{0}, r)} f(\mathbf{x}) \, d^n \mathbf{x} =: \frac{m(r)}{A_n}.$$

If we let f approach a Dirac's δ , we would end up with $m(r) = 1$ for all r . Thus, after integrating once more, we arrive at the solution $u = \Phi$, where $\Phi : \mathbb{R}^n \rightarrow \mathbb{R}$ is given by

$$\begin{aligned} \Phi(\mathbf{x}) &= \frac{-\ln(|\mathbf{x}|)}{2\pi} && \text{for } n = 2, \\ \Phi(\mathbf{x}) &= \frac{1}{(n-2)A_n|\mathbf{x}|^{n-2}} && \text{for } n \geq 3. \end{aligned}$$

For $n \geq 3$, this gives the unique radially symmetric solution which vanishes at infinity; for $n = 2$, however, no particular value of the dropped constant of integration distinguishes itself.

Now, put $m = 1$, replace $f(\mathbf{x})$ by $\varepsilon^{-n} f(\mathbf{x}/\varepsilon)$, and let $\varepsilon \rightarrow 0$. The resulting solution u will converge pointwise to Φ (except at $\mathbf{x} = \mathbf{0}$), while f becomes a Dirac δ in the limit. Thus we are tempted to conclude that

$$-\Delta\Phi = \delta$$

This is indeed true, but we first need to get into the theory of distributions in order to understand the rigorous meaning of the above equation.

We call Φ the *fundamental solution* for the Poisson equation. Even lacking the abstract theory, we can use it to solve the general Poisson equation.

Proposed solution to the Poisson equation:

$$u(\mathbf{x}) = \Phi * f(\mathbf{x}) = \int_{\mathbb{R}^n} \Phi(\mathbf{y})f(\mathbf{x} - \mathbf{y}) d^n \mathbf{y} \tag{9}$$

We expect this to solve the equation because formally, we get

$$\Delta u = \Delta(\Phi * f) = (\Delta\Phi) * f = \delta * f = f.$$

However, this calculation is hard to justify by elementary means, but we can do it via a detour.

Our main tool is Green's second identity. We often need to employ it on a region containing the singularity of Φ , which is of course not allowed. So the trick is to remove a ball centered on the singularity, use Green's second identity on the rest of the region, and then let the radius of the ball tend to zero. This leads to two limits handled by the following two lemmas.

Lemma 9. *If the function v is continuous near $\mathbf{0}$, then*

$$-\int_{\partial B(\mathbf{0},r)} v \partial_\nu \Phi dS = \int_{\partial B(\mathbf{0},r)} v dS \rightarrow v(\mathbf{0}) \quad \text{as } r \rightarrow 0.$$

Proof. The identity follows from the fact that the normal derivative on $\partial B(\mathbf{0}, r)$ is the radial derivative, with the constant value $\partial_\nu \Phi = -1/(A_n r^{n-1})$ on the sphere. And then the limit follows by continuity of v . ■

Lemma 10. *If the function v is C^1 near $\mathbf{0}$, then*

$$\int_{\partial B(\mathbf{0}, r)} \Phi \partial_\nu v \, dS \rightarrow 0 \quad \text{as } r \rightarrow 0.$$

Proof. The normal derivative $\partial_\nu v$ is really $\mathbf{v} \cdot \nabla v$, but \mathbf{v} is a unit vector, and ∇v is bounded near $\mathbf{0}$, since v is assumed to be C^1 . Because the integral of $|\Phi|$ tends to zero as $r \rightarrow 0$, the integral of the first term will vanish in the limit. ■

Theorem 11. *Assume that $f \in C_c^2(\mathbb{R}^n)$. Then the function u given by (9) is a solution to the Poisson equation (6).*

Proof. Thanks to translation invariance, we only need to prove that $-\Delta(\Phi * f)(\mathbf{0}) = f(\mathbf{0})$. (For any $\mathbf{x}_0 \in \mathbb{R}^n$, put $\tilde{f}(\mathbf{x}) = f(\mathbf{x}_0 + \mathbf{x})$. Then $\Phi * \tilde{f}(\mathbf{x}) = \Phi * f(\mathbf{x}_0 + \mathbf{x})$, so $-\Delta(\Phi * \tilde{f})(\mathbf{0}) = f(\mathbf{0})$ implies $-\Delta(\Phi * f)(\mathbf{x}) = f(\mathbf{x})$.)

First, we note that

$$\int_{\partial B(\mathbf{0}, r)} \Phi \, d^n \mathbf{x} = \begin{cases} -r \ln r & \text{for } n = 2, \\ r/(n-2) & \text{for } n \geq 3. \end{cases}$$

Integrating this with respect to r , we conclude that Φ is integrable (meaning the integral of $|\Phi|$ is finite) over $B(\mathbf{0}, r)$, and hence over any bounded subset of \mathbb{R}^n . From this, we conclude that not only is

$$u(\mathbf{x}) = \Phi * f(\mathbf{x}) = \int_{\mathbb{R}^n} \Phi(\mathbf{y}) f(\mathbf{x} - \mathbf{y}) \, d^n \mathbf{y}$$

well defined, but u is C^2 as well, and in fact

$$\Delta u = \Phi * \Delta f.$$

This is easy if f has compact support; it is also true if f and its derivatives up to second order vanish sufficiently fast at infinity, but we are not going to bother with this refinement.

In particular,

$$\Delta u(\mathbf{0}) = \int_{\mathbb{R}^n} \Phi(\mathbf{y}) \Delta f(-\mathbf{y}) \, d^n \mathbf{y} = \int_{\mathbb{R}^n} \Phi(\mathbf{y}) \Delta f(\mathbf{y}) \, d^n \mathbf{y}$$

(using the symmetry of Φ to get rid of the minus sign for convenience). We want to use Green's second identity to move the Laplacian to Φ instead. But then we need

to restrict attention to a bounded region, and we need to avoid the singularity of Φ at the origin. (It is this singularity that allows the answer to be non-zero, after all.) Pick R sufficiently large so $f(\mathbf{y}) = 0$ for $|\mathbf{y}| \geq R$, so that integrating over $B(\mathbf{0}, R)$ does not change the integral. Then, thanks to the integrability of Φ near the origin, we find

$$\Delta u(\mathbf{0}) = \lim_{r \rightarrow 0} \int_{B(\mathbf{0}, R) \setminus B(\mathbf{0}, r)} \Phi(\mathbf{y}) \Delta f(\mathbf{y}) \, d^n \mathbf{y}.$$

This integral can be transformed by Green’s second identity, resulting in

$$\Delta u(\mathbf{0}) = \lim_{r \rightarrow 0} \left(\int_{B(\mathbf{0}, R) \setminus B(\mathbf{0}, r)} \Delta \Phi(\mathbf{y}) f(\mathbf{y}) \, d^n \mathbf{y} + \int_{\partial(B(\mathbf{0}, R) \setminus B(\mathbf{0}, r))} (\Phi \partial_\nu f - f \partial_\nu \Phi) \, dS \right).$$

The first integral vanishes because $\Delta \Phi = 0$, and in the second integral we can ignore the outer boundary $\partial B(\mathbf{0}, R)$ because $f = \partial_\nu f = 0$ there. Finally, the normal vector ν points *inward* on the inner boundary $\partial B(\mathbf{0}, r)$, so we get a sign change when we consider the outward point normal instead. Thus we have

$$\Delta u(\mathbf{0}) = \lim_{r \rightarrow 0} \int_{\partial B(\mathbf{0}, r)} (f \partial_\nu \Phi - \Phi \partial_\nu f) \, dS.$$

The two preceding lemmas then yield $\Delta u(\mathbf{0}) = -f(\mathbf{0})$. ■

Bounded domains and Green’s function. Let Ω be a bounded domain with piecewise C^1 boundary. The weak maximum principle (even without the “nice” boundary) immediately shows that the Dirichlet problem

$$\begin{aligned} -\Delta u &= f && \text{in } \Omega, \\ u &= g && \text{on } \partial\Omega \end{aligned} \tag{DP}$$

has at most one solution for any given data f and g .

Our goal in this section is to generalize the representation formula (9) to this setting. We first concentrate on homogeneous boundary data, i.e., $g = 0$. It will turn out that we get the case for $g \neq 0$ “for free”.

First, rewrite (9) to read

$$u(\mathbf{x}) = \int_{\mathbb{R}^n} \Phi(\mathbf{x} - \mathbf{y}) f(\mathbf{y}) \, d^n \mathbf{y}.$$

A similar formula for solutions on Ω might look like

$$u(\mathbf{x}) = \int_{\Omega} G_{\mathbf{y}}(\mathbf{x}) f(\mathbf{y}) \, d^n \mathbf{y}$$

instead. To ensure that this satisfies $u = 0$ on $\partial\Omega$, we want $G_{\mathbf{y}}(\mathbf{x}) = 0$ for $\mathbf{x} \in \partial\Omega$. Apart from that, we want $G_{\mathbf{y}}(\mathbf{x})$ (as a function of \mathbf{x}) to be as much “like” $\Phi(\mathbf{x} - \mathbf{y})$ as possible. To make that precise:

Definition. Assume that, for each $\mathbf{y} \in \Omega$, there exists a function $H_{\mathbf{y}} \in C(\overline{\Omega})$ which is harmonic in Ω and satisfies

$$H_{\mathbf{y}}(\mathbf{x}) = \Phi(\mathbf{x} - \mathbf{y}) \quad \text{for } \mathbf{x} \in \partial\Omega.$$

By the maximum principle, this function is unique. Then the function $G_{\mathbf{y}}$ defined by

$$G_{\mathbf{y}}(\mathbf{x}) = \Phi(\mathbf{x} - \mathbf{y}) - H_{\mathbf{y}}(\mathbf{x})$$

is called the *Green’s function* associated with Ω . Clearly, this function is continuous on $\overline{\Omega} \setminus \{\mathbf{y}\}$ and harmonic on $\Omega \setminus \{\mathbf{y}\}$, and it vanishes on $\partial\Omega$.

To discover how this helps us write a solution formula for the Dirichlet problem, we concentrate first on the Φ part.

For simplicity, assume that $\mathbf{y} = \mathbf{0}$. If $\varepsilon > 0$ is small enough, $\overline{B(\mathbf{0}, \varepsilon)} \subset \Omega$. Put $\Omega_{\varepsilon} = \Omega \setminus \overline{B(\mathbf{0}, \varepsilon)}$, and apply Green’s second identity to Φ and an arbitrary function $u \in C^2(\overline{\Omega})$:

$$\begin{aligned} \int_{\Omega_{\varepsilon}} (u \Delta \Phi - \Phi \Delta u) d^n \mathbf{x} &= \int_{\partial\Omega_{\varepsilon}} (u \partial_{\nu} \Phi - \Phi \partial_{\nu} u) dS \\ &= \int_{\partial\Omega} (u \partial_{\nu} \Phi - \Phi \partial_{\nu} u) dS - \int_{\partial B(\mathbf{0}, \varepsilon)} (u \partial_{\nu} \Phi - \Phi \partial_{\nu} u) dS, \end{aligned}$$

where the minus sign in front of the last integral is due to the direction of the normal vector ν on $\partial B(\mathbf{0}, \varepsilon)$ pointing out of the ball, whereas the normal vector on that part of $\partial\Omega_{\varepsilon}$ points into the ball.

Now let $\varepsilon \rightarrow 0$. The red term on the left is already zero, while the boundary integral of the red term on the right will vanish in the limit by Lemma 10. Finally, the integral of the green term will converge to $-u(\mathbf{0})$ by Lemma 9, so we end up with the representation formula

$$u(\mathbf{y}) = - \int_{\Omega} \Phi \Delta u d^n \mathbf{x} - \int_{\partial\Omega} (u \partial_{\nu} \Phi - \Phi \partial_{\nu} u) dS, \quad \mathbf{y} \in \Omega. \quad (10)$$

(We proved it only for $\mathbf{y} = \mathbf{0}$, but translation invariance ensures that the result generalizes. Or you could rerun the proof with $\Phi(\mathbf{x})$ replaced by $\Phi_{\mathbf{y}}(\mathbf{x}) = \Phi(\mathbf{x} - \mathbf{y})$ and $B(\mathbf{0}, \varepsilon)$ by $B(\mathbf{y}, \varepsilon)$.)

Repeat the same calculation with Φ replaced by $H_{\mathbf{y}}$. This is much easier, since $H_{\mathbf{y}}$ is harmonic we do not need to remove a ball: Green’s second identity immediately yields

$$\int_{\Omega} (u \Delta H_{\mathbf{y}} - H_{\mathbf{y}} \Delta u) d^n \mathbf{x} = \int_{\partial\Omega} (u \partial_{\nu} H_{\mathbf{y}} - H_{\mathbf{y}} \partial_{\nu} u) dS,$$

where again the red term vanishes. Rearranging this into

$$\int_{\Omega} H_y \Delta u \, d^n \mathbf{x} + \int_{\partial\Omega} (u \partial_\nu H_y - H_y \partial_\nu u) \, dS = 0$$

and adding it to the right hand side of (10), we obtain

$$u(\mathbf{y}) = - \int_{\Omega} G_y \Delta u \, d^n \mathbf{x} - \int_{\partial\Omega} (u \partial_\nu G_y - G_y \partial_\nu u) \, dS$$

where the red term is zero by construction, so we finally have

$$u(\mathbf{y}) = - \int_{\Omega} G_y \Delta u \, d^n \mathbf{x} - \int_{\partial\Omega} u \partial_\nu G_y \, dS, \quad \mathbf{y} \in \Omega.$$

We have proved

Theorem 12. *If $\Omega \subset \mathbb{R}^n$ is a bounded region with C^1 boundary, and if Ω admits a Green's function G , then the solution $u \in C^2(\bar{\Omega})$ of (DP), if it exists, is given by*

$$u(\mathbf{y}) = \int_{\Omega} G_y(\mathbf{x}) f(\mathbf{x}) \, d^n \mathbf{x} - \int_{\partial\Omega} \partial_\nu G_y(\mathbf{x}) g(\mathbf{x}) \, dS(\mathbf{x}), \quad \mathbf{y} \in \Omega. \quad (11)$$

Notice the careful wording: We have not shown that the Dirichlet problem has a solution. We have only shown what it must be, if it solves the problem.

We now show that Green's function is symmetric.

Proposition 13. *Assume that Ω is a bounded region with piecewise C^1 boundary, and which admits a Green's function. Then the Green's function G for this region satisfies $G_y(\mathbf{x}) = G_x(\mathbf{y})$, whenever $\mathbf{x}, \mathbf{y} \in \Omega$.*

Proof. First a “physicists’ proof” to reveal the essentials: Since G_x vanishes on $\partial\Omega$ and $-\Delta G_x(\mathbf{z}) = \delta(\mathbf{z} - \mathbf{x})$, and similarly for G_y , Green's second identity yields

$$\begin{aligned} 0 &= \int_{\partial\Omega} (G_x \partial_\nu G_y - G_y \partial_\nu G_x) \, dS = \int_{\Omega} (G_x \Delta G_y - G_y \Delta G_x) \, d^n \mathbf{z} \\ &= \int_{\Omega} (-G_x(\mathbf{z}) \delta(\mathbf{z} - \mathbf{y}) + G_y(\mathbf{z}) \delta(\mathbf{z} - \mathbf{x})) \, d^n \mathbf{z} = -G_x(\mathbf{y}) + G_y(\mathbf{x}). \end{aligned}$$

Our problem with this proof is that Green's second identity, as we know it, requires C^2 functions, which is not the case here. So we remove some small balls around the troublesome points \mathbf{x} and \mathbf{y} : Put $\Omega_\varepsilon = \Omega \setminus (B(\mathbf{x}, \varepsilon) \cup B(\mathbf{y}, \varepsilon))$, where $\varepsilon > 0$ is small enough so that the two removed balls lie within Ω and have no point in common.

Now ΔG_x and ΔG_y are harmonic in Ω_ε , so Green's theorem is applicable:

$$\begin{aligned} 0 &= \int_{\Omega_\varepsilon} (G_x \Delta G_y - G_y \Delta G_x) d^n z = \int_{\partial \Omega_\varepsilon} (G_x \partial_\nu G_y - G_y \partial_\nu G_x) dS \\ &= - \left(\int_{\partial B(x, \varepsilon)} + \int_{\partial B(y, \varepsilon)} \right) (G_x \partial_\nu G_y - G_y \partial_\nu G_x) dS, \end{aligned}$$

where we dropped the integral over $\partial \Omega$, because the integrand is zero there. The minus sign is due to the reversed direction of the normal vector on the boundary of the balls versus the boundary of Ω_ε . We have already dealt with this sort of boundary integral in the proof of Theorem 12, so here we go again! Look at the first of the two boundary integrals above:

$$\int_{\partial B(x, \varepsilon)} (G_x \partial_\nu G_y - G_y \partial_\nu G_x) dS.$$

The integral of the red term is zero, for G_x is constant on $\partial B(x, \varepsilon)$, and

$$\int_{\partial B(x, \varepsilon)} \partial_\nu G_y dS = \int_{B(x, \varepsilon)} \Delta G_y d^n z = 0$$

by the divergence theorem. Even without harmonicity, this would vanish in the limit as $\varepsilon \rightarrow 0$.

For the green term, note that $-\partial_\nu G_x = 1/(A_n \varepsilon^{n-1})$ on $\partial B(x, \varepsilon)$, so together with the above calculation we get

$$\int_{\partial B(x, \varepsilon)} (G_x \partial_\nu G_y - G_y \partial_\nu G_x) dS = \int_{\partial B(x, \varepsilon)} G_y dS \rightarrow G_y(\mathbf{x}) \quad \text{as } \varepsilon \rightarrow 0.$$

Similarly,

$$\int_{\partial B(y, \varepsilon)} (G_x \partial_\nu G_y - G_y \partial_\nu G_x) dS = \int_{\partial B(y, \varepsilon)} G_x dS \rightarrow G_x(\mathbf{y}) \quad \text{as } \varepsilon \rightarrow 0,$$

and the proof is complete. ■

Remark. A different take the proof of symmetry of Green's function:

Let $f, g \in C(\overline{\Omega})$, and let u, v solve $-\Delta u = f$ and $-\Delta v = g$ in Ω , $u = v = 0$ on $\partial \Omega$. Thus

$$u(\mathbf{y}) = \int_{\Omega} G_y(\mathbf{x}) f(\mathbf{x}) d^n \mathbf{x}, \quad v(\mathbf{y}) = \int_{\Omega} G_y(\mathbf{x}) g(\mathbf{x}) d^n \mathbf{x}.$$

Since u and v vanish on $\partial \Omega$, Green's second identity applied to u and v implies

$$\int_{\Omega} u(\mathbf{y}) g(\mathbf{y}) d^n \mathbf{y} = \int_{\Omega} v(\mathbf{y}) f(\mathbf{y}) d^n \mathbf{y}.$$

Plugging in the above formulas for u and v , swapping the integration variables x and y in one of the integrals, and rearranging, we get

$$\int_{\Omega} \int_{\Omega} (G_y(x) - G_x(y))f(x)g(y) d^n x d^n y = 0,$$

from which the desired conclusion quickly follows. (Take f and g to be approximate delta functions concentrated near two different points in Ω .)

It may be interesting to study the above argument from an elementary linear algebra point of view. The last part resembles the proof that, if A is a quadratic matrix satisfying $y^T A x = x^T A y$ for all vectors x and y , then A is symmetric. (Just take $x = e_i$ and $y = e_j$.) The first part resembles the proof that the inverse of an invertible symmetric matrix is symmetric. Think of the “matrix” $G_x(y)$ (where x and y play the rôles of matrix indices) as the inverse of the operator $-\Delta$. And the symmetry of this operator is just the identity $\int_{\Omega} u \Delta v = \int_{\Omega} v \Delta u$. (Green’s second identity, when both functions vanish on the boundary.)

Green’s function for balls. Here we compute Green’s function for the open unit ball B^n in \mathbb{R}^n . I will drop some exponents and write B instead of B^m and $S = \partial B$ instead of S^{n-1} .

If $y \in B$, we need to find a harmonic function on B with the same values as $\Phi(x - y)$ for $x \in S$. The trick is to “put a charge” at a suitable point w outside B . It turns out that $w = y/|y|^2$ does the trick. Since $|x| = 1$, we find

$$|x - y|^2 = 1 + |y|^2 - 2x \cdot y$$

and

$$|x - w|^2 = 1 + |w|^2 - 2x \cdot w = 1 + \frac{1}{|y|^2} - 2 \frac{x \cdot y}{|y|^2} = \frac{|y - x|^2}{|y|^2},$$

so that

$$|w - x| = \frac{|y - x|}{|y|}.$$

For $n \geq 3$, this gives

$$\Phi(w - x) = \frac{1}{(n - 2)A_n |w - x|^{n-2}} = |y|^{n-2} \Phi(y - x),$$

so we should put

$$H_y(x) = \frac{\Phi(w - x)}{|y|^{n-2}} = \frac{1}{(n - 2)A_n |y|^{n-2} |w - x|^{n-2}}$$

where we note that

$$|y|^2|w - x|^2 = |y|^2(|w|^2 + |x|^2 - 2x \cdot w) = 1 + |x|^2|y|^2 - 2x \cdot y,$$

so we get

$$G_y(x) = \frac{1}{(n-2)A_n} \left(\frac{1}{|x-y|^{n-2}} - \frac{1}{(1+|x|^2|y|^2-2x \cdot y)^{(n-2)/2}} \right) \quad (n \geq 3)$$

which is indeed symmetric in x, y as expected.

Next, the same computation for $n = 2$: We find (when $x \in \mathbb{S}$)

$$\Phi(w - x) = -\frac{\ln|w - x|}{2\pi} = \frac{\ln|y|}{2\pi} + \Phi(y - x),$$

so we put

$$H_y(x) = -\frac{\ln|y|}{2\pi} + \Phi(w - x) = -\frac{\ln(|y||w - x|)}{2\pi}$$

and get

$$G_y(x) = -\frac{\ln|y - x|}{2\pi} + \frac{\ln(1 + |x|^2|y|^2 - 2x \cdot y)}{4\pi} \quad (n = 2)$$

which is again symmetric. We can also write this on the form

$$G_y(x) = -\frac{\ln(|x|^2 + |y|^2 - 2x \cdot y)}{4\pi} + \frac{\ln(1 + |x|^2|y|^2 - 2x \cdot y)}{4\pi} \quad (n = 2)$$

When solving the Dirichlet problem on the unit disk $\mathbb{D} \subset \mathbb{R}^2$, we need the normal derivative $\partial_\nu G_y(x)$ for $x \in \mathbb{S}$ which is simply the derivative of $G_y(tx)$ taken at $t = 1$, resulting in

$$\partial_\nu G_y(x) = \frac{-2 + 2x \cdot y + 2|y|^2 - 2x \cdot y}{4\pi(1 + |y|^2 - 2x \cdot y)} = \frac{|y|^2 - 1}{2\pi(1 + |y|^2 - 2x \cdot y)}.$$

If we put $r = |y|$ and let θ be the angle between the two vectors x and y , we get

$$-\partial_\nu G_y(x) = \frac{1}{2\pi} \frac{1 - r^2}{1 + r^2 - 2r \cos \theta}$$

which is known as the *Poisson kernel*. Referring back to (11), the solution to the Dirichlet problem $-\Delta u = 0$ on \mathbb{D} with $u(\cos \theta, \sin \theta) = g(\theta)$ should then be given by

$$u(r \cos \varphi, r \sin \varphi) = \frac{1 - r^2}{2\pi} \int_{-\pi}^{\pi} \frac{g(\varphi - \theta)}{1 + r^2 - 2r \cos \theta} d\theta$$

This is known as *Poisson's integral formula*. It was originally derived using separation of variables and a Fourier analysis:

For this, we may note that the Laplace operator in polar coordinates takes the form

$$\Delta u = r^{-1}(ru_r)_r + r^{-2}u_{\theta\theta}.$$

We look for harmonic functions of the form $u = R(r)\Theta(\theta)$, resulting in $r^{-1}(rR')'\Theta + r^{-2}R\Theta'' = 0$. After separating the variables, we are left with $\Theta'' = -\lambda\Theta$ and $(rR')' = \lambda r^{-1}R$. Since Θ must be 2π -periodic, we must put $\lambda = n^2$ for an integer n , so the R equation is $r(rR')' = n^2R$. We try $R = r^k$, and get $rR' = kr^k$, $r(rR')' = k^2r^k$, with the non-trivial solutions $k = \pm n$. Rejecting solutions with negative k (which have a singularity at $r = 0$), we are therefore left with $R = r^{|n|}$. Thus we are led to look for solutions of the form

$$u(r \cos \varphi, r \sin \varphi) = \sum_{n=-\infty}^{\infty} c_n r^{|n|} e^{in\varphi}.$$

Matching this to $g(\theta)$ at $r = 1$, we should have

$$c_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} g(\theta) e^{-in\theta} d\theta,$$

and therefore

$$u(r \cos \varphi, r \sin \varphi) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sum_{n=-\infty}^{\infty} g(\theta) r^{|n|} e^{in(\varphi-\theta)} d\theta = \frac{1}{2\pi} \int_{-\pi}^{\pi} g(\theta) P(r, \varphi - \theta) d\theta,$$

where

$$P(r, \theta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} r^{|n|} e^{in\theta}.$$

Here we note that

$$\sum_{k=0}^{\infty} r^k e^{\pm ik\theta} = \frac{1}{1 - r e^{\pm ik\theta}},$$

so

$$\begin{aligned} P(r, \theta) &= \frac{1}{2\pi} \left(\frac{1}{1 - re^{ik\theta}} + \frac{1}{1 - re^{-ik\theta}} - 1 \right) \\ &= \frac{1}{2\pi} \frac{1 - r^2}{1 + r^2 - 2r \cos \theta}, \end{aligned}$$

which we recognize as the Poisson kernel introduced above.