

First order quasilinear equations

Harald Hanche-Olsen

More general quasilinear equations

Here we look at the general quasilinear equation in two dimensions:

$$au_x + bu_y = c, \quad (1)$$

in which a , b , and c are *really* $a(x, y, u(x, y))$, $b(x, y, u(x, y))$, and $c(x, y, u(x, y))$. (It is the dependence of the coefficients on u that makes the equations *quasilinear*.)

Consider any smooth curve $(x(\tau), y(\tau))$. Then, assuming u is a classical solution of (1), we put $z(\tau) = u(x(\tau), y(\tau))$, and find

$$z'(\tau) = x'(\tau)u_x + y'(\tau)u_y$$

so that if $x'(\tau) = a$ and $y'(\tau) = b$, then

$$z' = au_x + bu_y = c.$$

This all means that (x, y, z) satisfy the *characteristic equations*

$$x'(\tau) = a(x, y, z), \quad y'(\tau) = b(x, y, z), \quad z'(\tau) = c(x, y, z). \quad (2)$$

To summarize so far: *Assume that u is a classical solution of (1). Through each point in the graph of u ,*

$$\text{graph}(u) = \{(x, y, z) \mid z = u(x, y)\},$$

there passes a characteristic curve $(x(\tau), y(\tau), z(\tau))$ solving (2). Moreover, each such characteristic curve will lie within the graph of u .

The solution strategy for (1) can now be explained: Since the graph of a classical solution is a union of characteristic curves, we try to construct solutions by putting together characteristic curves.

The graph of u will be two-dimensional, and a characteristic curve is one-dimensional; so it makes sense to use another variable to keep track of the characteristic curves.

In other words, to construct a solution, we look for three functions $x(\sigma, \tau)$, $y(\sigma, \tau)$, $z(\sigma, \tau)$ which parametrize a characteristic curve as function of τ for each σ . In other words, they should satisfy

$$x_\tau = a, \quad y_\tau = b, \quad z_\tau = c$$

where, as always, a , b , and c are considered functions of x , y , and z .¹ Additionally, we assume that these are C^1 functions, and that $(x(\sigma, \tau), y(\sigma, \tau))$ has a C^1 inverse, mapping (x, y) to (σ, τ) . Then we can define u by

$$u(x(\sigma, \tau), y(\sigma, \tau)) = z(\sigma, \tau).$$

Differentiating this equation with respect to τ yields $x_\tau u_x + y_\tau u_y = z_\tau$, which is the same as (1).

To make this construction more concrete, putting $\tau = 0$ yields a parametric curve γ in the (x, y) -plane: $(x(\sigma, 0), y(\sigma, 0))$, or, if we add a coordinate, a curve Γ in $\text{graph}(u)$ parametrized as $(x(\sigma, 0), y(\sigma, 0), z(\sigma, 0))$. The requirement that $(x(\sigma, \tau), y(\sigma, \tau))$ has a C^1 inverse implies that the matrix $\begin{pmatrix} x_\sigma & x_\tau \\ y_\sigma & y_\tau \end{pmatrix}$ is non-singular. At $\tau = 0$ this simply means that Γ is not tangent to the characteristic curves. We refer to this as the *non-characteristic condition*.

We now see what is a natural condition to impose in order to obtain a unique solution to (1): Namely, given a curve γ in the (x, y) -plane, and a function g on γ , assume that the curve Γ given by points (ξ, η, ζ) with (ξ, η) on γ and $\zeta = g(\xi, \eta)$ satisfies the non-characteristic condition. Then (1) has a solution u satisfying the condition

$$u = g \text{ on } \gamma. \tag{3}$$

This solution will exist on a neighbourhood of γ , and be unique there. (Though we must be careful with any end points of γ : They should not be points on γ themselves, as we would then be obliged to extend the solution beyond the end of γ .)

A problem of the form (1) with “initial” data (3) is called a *Cauchy problem* for (1). More generally, a Cauchy problem for a PDE is the problem of solving the PDE along with certain conditions along a curve, or more generally a hypersurface. Often, as is the case here, the PDE more or less dictates the proper form of the Cauchy problem after some analysis.

¹Well, they were functions of x , y , and u , right? But in this solution, we should think of u and z as being the same. We write u when we emphasize the solution $u(\tau, x)$, but z when we think of the characteristic curves, as in $z(\tau)$ or $z(\sigma, \tau)$. After you have gained some experience, you may find it easier to forget about z and just write u . But this may be too confusing in the beginning.

The general quasilinear equation in higher dimensions

We can do the exact same procedure, with the same arguments, in n dimensions as in the 2-dimensional case. Here we summarize the construction *very* briefly.

A general quasilinear equation then takes the form

$$\mathbf{a}(\mathbf{x}, u(\mathbf{x})) \cdot \nabla u(\mathbf{x}) = c(\mathbf{x}, u(\mathbf{x})), \quad \mathbf{x} \in \mathbb{R}^n, \quad (4)$$

with given functions \mathbf{a} and c .

The *characteristic equations* become

$$\mathbf{x}'(\tau) = \mathbf{a}(\mathbf{x}(\tau), z(\tau)), \quad z'(\tau) = c(\mathbf{x}(\tau), z(\tau)),$$

or written in a more compact form:

$$\mathbf{x}' = \mathbf{a}(\mathbf{x}, z), \quad z' = c(\mathbf{x}, z). \quad (5)$$

Assume now that we are given the PDE (4) with the extra condition

$$u(\xi) = g(\xi), \quad \xi \in \gamma, \quad (6)$$

where $\gamma \subset \mathbb{R}^n$ is a *hypersurface*, i.e., a surface of dimension $n - 1$. The non-characteristic condition now says that $\mathbf{a}(\xi, g(\xi))$ is not tangent to γ for any $\xi \in \gamma$.

For any $\xi \in \gamma$, standard ODE theory guarantees the existence of a solution of (5) satisfying $\mathbf{x}(0) = \xi$ and $z(0) = g(\xi)$. Write $(\mathbf{x}(\tau, \xi), z(\tau, \xi))$ for this solution, and define

$$u(\mathbf{x}(\tau, \xi)) = z(\mathbf{x}(\tau, \xi)), \quad \xi \in \gamma,$$

where again, we can show that this is well defined (for τ sufficiently close to 0) by using the inverse function theorem. Further, the $n + 1$ variables t, ξ are essentially only n variables, because γ is $(n - 1)$ -dimensional. The proof that this produces a classical solution is similar to the 2-dimensional case.